

DAI Jingen, WANG Chengshan, Zhu Dicheng, POLAT Ali and LI Yalin, 2015. Subduction Initiation and Generation of the Xigaze Ophiolite, Southern Tibet. *Acta Geologica Sinica* (English Edition), 89(supp. 2): 5.

Subduction Initiation and Generation of the Xigaze Ophiolite, Southern Tibet

DAI Jingen^{1,*}, WANG Chengshan¹, Zhu Dicheng¹, POLAT Ali² and LI Yalin¹

1 *State Key Laboratory of Biogeology and Environmental Geology, School of Earth Sciences and Resources, Research Center for Tibetan Plateau Geology, China University of Geosciences (Beijing), Beijing 100083, China*

2 *Department of Earth and Environmental Sciences, University of Windsor, Windsor, ON, Canada N9B 3P4*

The Cretaceous Xigaze ophiolite is best exposed at the central Yarlung Zangbo Suture Zone (YZSZ, Tibet) which also includes the Gangdese arc and the Xigaze forearc basin. This study reports new geochronological and geochemical data for this ophiolite to revisit its geodynamic and petrogenetic evolution. The Xigaze peridotites have low CaO and Al₂O₃ contents and U-shaped Rare Earth Element (REE) patterns, suggesting that they are residues after moderate to high degrees of partial melting and were modified by infiltration of Light Rare Earth Element (LREE)-enriched boninitic melts. The Xigaze crustal rocks belong to two groups: Mid-Ocean-Ridge-Basalt (MORB)-like rocks and boninitic rocks showing a uniform LREE depletion and flat to LREE enrichment on chondrite-normalized patterns, respectively. Geochemically, both groups show the influence of

subducting oceanic slab-derived fluids. LA-ICPMS zircon U-Pb and Lu-Hf analyses from dolerite and quartz diorite dikes, which intruded into the mantle peridotite, and dolerite sheeted sills show that they were generated between 127–124 Ma. The zircons possess positive $\epsilon_{\text{Hf}}(t)$ values ranging from +7.5 to +17.3. Taking into account the geological and geochronological characteristics of the central-western YZSZ, we propose that ophiolites in this region formed in a forearc spreading setting through rapid slab rollback during subduction initiation between 130–120 Ma. Following this stage of spreading, the forearc was stabilized and the zone of melting migrated beneath the Gangdese arc producing the voluminous Late Cretaceous granitoids displaying depleted mantle-type Hf isotopic compositions. Our model provides a new explanation for the generation and evolution of forearc-type ophiolites.

* Corresponding author. E-mail: djgtibet@163.com