

LI Xuping, KONG Fanmei, CHEN Hongkai and ZHANG Xin, 2015. Rodingite in the Purang Ophiolite and Its Geological Implication, Southwest Tibet. *Acta Geologica Sinica* (English Edition), 89(supp.): 41-42.

## Rodingite in the Purang Ophiolite and Its Geological Implication, Southwest Tibet

LI Xuping<sup>\*</sup>, KONG Fanmei, CHEN Hongkai and ZHANG Xin

*Shandong University of Science and Technology, Qingdao 266590, China*

The Yarlung Zangbo Suture Zone (YZSZ) separates the Lhasa block of the Eurasian plate to the north from the Indian plate to the south, along a ~2000km discontinuous E–W trending suture zone, several heterogeneous remnants of the Neo-Tethyan oceanic floor have been preserved from the subduction recycling. The Purang ophiolite massif is one of the most spectacular exposures of western extension along the Indus-Yarlung Zangbo Suture between the Himalayan passive margin and the Gangdise active margin, which is a mantle thrust sheet of more than 650 km<sup>2</sup>, extending NWW in the western Tibet (Nicolas et al., 1981; Yang et al., 2011).

The Purang ophiolite consists mainly of spinel peridotites, including serpentized harzburgite, minor lherzolite and chromite-bearing dunite associated with small amounts of olivine pyroxenite. Gabbro and diabase dike were observed intruding and cutting through spinel peridotites. Along the northern margin there is a mélange of serpentized harzburgite, tectonically overlying upper Triassic-Cretaceous clastic rocks. Partly serpentized harzburgite accounts for more than 80% of the ultramafic rocks. Lherzolite, dunite and olivine pyroxenite mostly occur as irregular blocks or bands within the harzburgite, some of which have clear boundaries with the host (Li et al., 2015).

Rodingites occur as lens, block or vein along the fracture in the south margin of the Purang ultramafic massif, and it is associated with relatively intense serpentized peridotite. Since the Purang ultramafic rocks suffered only weak to moderate serpentization, rodingite only experienced early stage mineral assemblage of rodingitization (Li et al., 2007, 2008). Rodingite has intense crack texture and consists of amphibole, clinzoisite and chlorite. Accessory minerals are titanite and ilmenite. No relict primary minerals were found. Two generations of amphibole were found, the early one has edenitic composition, while the late amphiboles are Mg-hornblende, actinolite to tremolite, formed during rodingitization. Epidote-group minerals are clinzoisites

and chlorite is clinocllore in composition.

Rodingites from the Purang area contain average SiO<sub>2</sub> ~48.65%, CaO ~ 9.89%, TFe<sub>2</sub>O<sub>3</sub> ~4.34%, Na<sub>2</sub>O + K<sub>2</sub>O < 2%. Additionally, they have extraordinarily high MgO ~23.92% and low TiO<sub>2</sub> < 0.02% in average. Bulk-rock recalculation indicates the protolith of rodingite is a dark olivine gabbro. Loss of ignition (LOI) varies from 1.83–3.23 wt%, reflecting weekly degrees of rodingitization.

The chondrite normalized composition of the rodingite generally displays a slightly depleted LREE pattern. The multi-element diagram normalized to primary mantle presents that all samples are enriched in U, Pb and Sr anomalies, but almost depleted in other trace elements with negative anomalies in part high-field strength elements (e.g. Nb, Ti, Y). The  $\Sigma$ REE of rodingite is much lower than that in the gabbroic diabase, but higher than that of spinel-bearing ultramafic rocks in the Pruang massif.

As conclusions, the Purang rodingite shows a mantle source protolith. Some rodingites are located in the IAT domain in the Th/Yb versus Nd/Yb diagram, whereas others are in between N-MMORB and E-MORB domain. Positive anomalies in U, Pb and Sr indicate a subduction-related affinity. REE and trace elements of rodingites are all indicative of fore-arc geochemical characteristics based on the diagrams of V-Ti, La/Nb-Y and Zr/Y-Zr (The current study was supported by the Chinese Geological Survey Project (12120114061801) and the SGSTSP of the Chinese Academy of Sciences XDB03010201).

### References

- Li, X.P., Chen, H.K., Wang, Z.L., Wang, L.J., Yang, J.S., and Robinson, P., 2015. Textural evolution of spinel peridotite and olivine websterite in the Purang ophiolite complex, western Tibet. *Journal of Asian Earth Sciences*, 110: 55–71.
- Li, X.P., Rahn, M., and Bucher, K., 2008. Eclogite-facies metarodingites: phase relations in the system SiO<sub>2</sub>-Al<sub>2</sub>O<sub>3</sub>-FeO-Fe<sub>2</sub>O<sub>3</sub>-MgO-CaO-CO<sub>2</sub>-H<sub>2</sub>O, an example from the Zermatt-Saas Ophiolite. *Journal of Metamorphic Geology*, 26: 347–364.
- Li, X.P., Zhang, L.F., Wei, C.J., Ai, Y.L., and Chen, J., 2007. Petrology of rodingite derived from eclogite in western

\* Corresponding author. E-mail: lixuping@sdust.edu.cn

- Tianshan, China. *Journal of Metamorphic Geology*, 25: 363–382.
- Nicolas, A., Girardeau, J., Marcoux, J., Dupre, B., Wang, X., and Cao, Y., et al., 1981, The Xigaze ophiolite (Tibet): a peculiar oceanic lithosphere. *Nature*, 294: 414–417.
- Yang, J.S., Xiong, F.H., Guo, G.L., Liu, F., Liang, F.H., Chen, S.Y., Li, Z.L., and Zhang, L.W., 2011. The Dongbo ultramafic massif: a mantle peridotite in the western part of the Yarlung Zangbo suture zone, Tibet, with excellent prospects for a major chromite deposit. *Acta Petrol. Sinica*, 27: 3207–3222 (in Chinese with an English abstract).