

HOU Guiting, 2016. Mechanics of Mafic Dyke Swarms and Breakup of Supercontinent. *Acta Geologica Sinica* (English Edition), 90 (supp. 1): 155-156.

Mechanics of Mafic Dyke Swarms and Breakup of Supercontinent

HOU Guiting

*The Key Laboratory of Orogenic Belts and Crustal Evolution, Education Ministration,
School of Earth and Space Sciences, Peking University, Beijing 100871, China*

Dyke swarms can be divided into three types: parallel dyke swarms, radiating dyke swarms and fan-shape dyke swarm, for which the mechanisms of formation are different (Fig. 1). Parallel dyke swarms form in response to a regional stress field, for example, the NNW-trending mafic dyke swarms widespread developed in the the North China Craton (Hou et al., 2006a,b; Peng et al., 2007, 2011). The radiating dyke swarms are developed due to stress concentrations around a volcanic edifice, as in the case of the radiating dyke swarm at Spanish Peak, USA. The fan-shape dyke swarms conclude two parts: radiating dyke swarm and parallel dyke swarm, for example, Mackenzie dyke swarm (Halls, 1987; Ernst et al., 1995). In the factors controlling the emplacement of dyke swarm, fractures that formed prior to magmatism may play a role in dictating dyke swarm geometry. In most cases pre-existing fractures are induced by tectonic stresses and not by magma injection though magma injection can increase fracture size by fracture propagation at the dyke tip.

The regional stress field is very important factor to control the distribution of parallel dyke swarm. Usually the parallel dyke swarms are controlled by the regional

extension stress field which tectonic forces come from the boundaries of the plate. The “pre-existing hole” model (Fig. 2A) for the formation of radiating dyke swarms can explain the mechanism of small-scale radiating dyke swarms related to the volcanic edifice, but not provide an explanation for the size of giant radiating dyke swarms related to mantle plume (Park, 1995). In the case of the 2100 km long Mackenzie dyke swarm of the Canadian Shield, radiating dykes occur around a focal source but with distance, dyke orientations swing from N-S to NW-SE and the dyke swarm adopts the geometry of a parallel dyke swarm.

The mechanics of the giant radiating dyke swarms, i.e., Mackenzie dyke swarm, is very complex. A “Plug” model based on the results of two dimension finite element modeling is proposed to explain the mechanism of formation of the Mackenzie dyke swarm (Hou et al., 2010) (Fig. 2B). Hou’s “Plug” model can explain the development of radiating dykes concentrated around the focal area and parallel dykes far from the magma source during opening of the Grenville Ocean. The model is a “plug-like” model that invokes local stress concentrations




Fig. 1. Three types of dyke swarms. Arrow indicates the regional maximum principle compressive stress.

* Corresponding author. E-mail: gthou@pku.edu.cn




Fig. 2. Two mechanics models for the radiating dyke swarms. Model A—mechanics for small scale radiating dyke swarms related to volcanic edifice; Model B—mechanics for giant fan-shape dyke swarm related to mantle plume and ocean ridge.

around the magmatic focal area (boundary of Coppermine Lava) and a regional stress field far from the focal area, induced by opening of the, Grenville Ocean (ridge push) (Fig. 2B).

Since giant radiating dyke swarms and rifts developed in most of the cratons on Earth during the 1.3-1.2Ga periods, it can be suggested that these cratons were connected together in supercontinent Columbia, and that a supercontinent-wide extension event initiated its final breakup in the Late Mesoproterozoic (Zhai et al, 2003; Zhao et al., 2004; Hou et al., 2008a,b). After the final global breakup, Columbia was fragmented into separated continental blocks during 1.2-1.3 Ga. The giant radiating Mackenzie dyke swarm of 1.2-1.3 Ga is related to a super mantle plume below the Canadian Shield, and suggests that the upwelling of mantle plume and Grenville Ocean spreading both contribute to the final breakup of Columbia supercontinent.

References

- Ernst, R., Head, J., and Parfitt, E., 1995. Giant radiating dyke swarms on Earth and Venus. *Earth-Science Reviews*, 39: 1-58.
- Halls, H., 1987. Dyke swarms and continental rifting some concluding remarks. In: Halls, H.C., and Fahrig, W.F. (eds.), *Mafic Dyke Swarms*. Geological Association of Canada Special Paper, 134: 5-24.
- Hou, G., Kusky, T.M., Wang, C., and Wang, Y., 2010. Mechanics of the giant radiating Mackenzie dyke swarm: A paleostress field modeling. *Journal of Geophysical Research*, 115, B 02402: 1-14.
- Hou, G., Liu, Y., and Li, J., 2006a. The evidence of ~1.8 Ga extension of NCC from the U-Pb SHRIMP dating of mafic dyke in the Shandong Province. *Journal of Asian Earth Sciences*, 27: 392-401.
- Hou, G., Wang, C., and Li, J., 2006b. The Paleoproterozoic extension and reconstruction of paleostress field of the North China Craton. *Tectonophysics*, 422: 87-98.
- Hou, G., Santosh, M., and Qian, X., 2008a. Configuration of the Late Paleoproterozoic supercontinent Columbia: Insights from radiating mafic dyke swarms. *Gondwana Research*, 14: 395-409.
- Hou, G., Santosh, M., and Qian, X., 2008b. Tectonic constraints on 1.3~1.2 Ga final breakup of Columbia supercontinent from a giant radiating dyke swarm. *Gondwana Research*, 14: 561-566.
- Li, Z., Li, X., and Kinny, P., 1999. The breakup of Rodinia: did it start with a mantle plume beneath South China? *Earth and Planetary Science Letters*, 173: 171-181.
- Park, J., Buchan, K., and Harlan, S., 1995. A proposed giant radiating dyke swarm fragmented by the separation of Laurentia and Australia based on paleomagnetism of ca. 780 Ma mafic intrusions in western North America. *Earth and Planetary Science Letters*, 132: 129-139.
- Peng, P., Wouter, B., and Richard, E., 2011. U-Pb baddeleyite ages, distribution and geochemistry of 925 Ma mafic dykes and 900 Ma sills in the NCC: Evidence for a Neoproterozoic mantle plume. *Lithos*, 127: 210-221.
- Peng, P., Zhai, M., and Guo, J., 2007. Nature of mantle source contributions and crystal differentiation in the petrogenesis of the 1.78 Ga mafic dykes in the central North China Craton. *Gondwana Research*, 12(1-2): 29-46.
- Zhai, M., and Liu, W., 2003. Paleoproterozoic tectonic history of the North China Craton: a review. *Precambrian Research*, 122: 183-199.
- Zhao, G., Sun, M., and Wilde, S., 2004. A Paleo-Mesoproterozoic supercontinent: assembly, growth and breakup. *Earth-Science Reviews*, 67: 91-123.