

HAO Guibao, FANG Weixuan, GUO Yuqian and CAO Jingwei, 2017. Studies on Characteristics of Fracture Lithofacies in Wenxipo W-Cs-Rb Polymetallic Deposit, Hainan Province, China. *Acta Geologica Sinica* (English Edition), 91(supp. 1): 212-213.

## Studies on Characteristics of Fracture Lithofacies in Wenxipo W-Cs-Rb Polymetallic Deposit, Hainan Province, China

HAO Guibao<sup>1,2</sup>, FANG Weixuan<sup>1,2</sup>, GUO Yuqian<sup>2</sup> and CAO Jingwei<sup>3</sup>

1 *China University of Geosciences, Beijing 100083;*

2 *China Non-ferrous Metals Resource Geological Survey, Beijing 100012;*

3 *Hainan Danzhou Mining co., LTD.*

### 1 Introduction

Wenxipo W-Cs-Rb polymetallic deposits are located in Fengshou rubidium-cesium polymetallic prospecting area in Danzhou City, Hainan Province, China. The hornstone facies, a thermal metamorphic facies, developed around the diorite. Alteration type of wall rock including mainly potassium-rich alteration, silicification, and hornstone facies as well as chloritization, biotitization, tourmalinization, pyritization, and pyrrhotiteization. The relationships between the fracture and the mineralization regularity of tungsten were discussed in this study. The petrographic study with the statistics and fracture measuring, fissure fillings, suggested that fracture's angle in this study area might be divided into four groups, i.e., 9°, 29°, 47°, 69°, respectively. The scheelite mainly occurred in the interlayer-fracture, the inclination of scheelite stringer is about 45°, which is roughly same as stratigraphic tectonic deformation foliation, and part of scheelite stringer occurred in strata penetration fissure in the group. In the scheelite orebody, the more fracture density, the more fissure permeability, indicated that the fracture in the rocks might provide the migration channels for the ore-forming fluid and ore-contained space for the scheelite mineralization. Crashed zone and scheelite orebody at the higher ore-grade may be controlled by biotite diorite intrusion obviously. Indeed, Cataclastic lithofacies is proportional to the fracture density near the biotite diorite intrusion, fracture density up to 35 / m on average. Filling types and characteristics of fracture showed that the fracture fillings have formed in multi-phase features, fissure fillings included high temperature to low temperature mineral. The microscopic identification results show that the enrichment of scheelite mainly related to

tourmalinization and silicification closely, speculated that the scheelite mineralization might be mainly related to the magmatic hydrothermal mineralization of biotite diorite. The discovery of tungsten deposit has a large prospecting potential, however, the relationship between the fracture and W-Cs-Rb enrichment regularity in the study needs to be studied.

Rock fractures, including joint, cleavage and fissure, which has a very close relationship for formations of metal ore deposit (such as Yan et al., 2015; Fang et al., 2001; Liao et al., 2013; Wang et al., 2016; Wang et al., 2016; Lin et al., 2012). Rubidium-cesium polymetallic deposit belongs to a skarn-greisen-type, which were found by Marine Geological Survey Institute of Hainan Province. However, previous study paid little attention to fracture lithofacies.

China Non-ferrous Metals Resource Geological Survey has found and delineated the tungsten orebody, using comprehensive research, comprehensive assessment, comprehensive prospecting and evaluation. It was believed that tungsten orebody is main mineral commodity, Cs and Rb is associated mineral commodity. Occurrence state of tungsten is scheelite and wolframite, they are deemed to be available industrial minerals. However, Occurrence state of Cs and Rb are in biotite and phlogopite, which may be the Cs-Rb concentration minerals for industrial utilization.

### 2 Methodology

Fractures in the drilling-core may provide direct information for the study on the fracture lithofacies. For it may provide quantitative and semi-quantitative description of the varied characteristics of the fissure development in the study area. The main research

**Table1. DZK1097 Fracture permeability characteristics of table in the drill core**

Research unit	Depth	Crack density( strip/cm)	Width of fissure (cm)/d	Permeability (cm <sup>2</sup> )/10 <sup>-4</sup>
diorite	0-149.54m	36.48	0.19	4.53
hornstone	149.54-296.43m	52.10	0.12	0.90
diorite	296.43-395.83m	25.92	0.14	0.67
hornstone	395.83-449.72m	18.09	0.15	0.98
diorite	449.72-481.44m	3.50	0.20	0.23
hornstone	481.44-488.61m	6.67	0.13	0.44

methods in this study are as follows: (1) to divide the type of fractures in the core library. (2) to measure for the density and inclination of fissures. (3) to measure the filling and opening degree of fissure. (4) the mechanism coupling relationships between fracture and ore-forming fluid maybe indicated by the permeability of rock fissure. (5) to observe the micro-fractures in office study including the observation and identification of micro-structures, the statistics of micro-fractures, the intertwined relationship of micro-fractures, the quantitative description of the micro-fractures development of the core and the contribution of the micro-fractures to the reservoir structure or reservoir. (6) To study on fracture fillings combining the macroscopic observation in field with office research methods. We have completed drilling-core logging by fluorescent scanning method in the field in order to get the formations on the fracture fillings such as scheelite, fluorite and calcite with REE and to summary on the enrichment and mineralization regularity of scheelite. The filling-minerals in the micro-fractures were observed by EPMA. W-Cs-Rb hosted minerals filling in fractures in the study area are detail studied by EPMA and XRD in the office studies.

### 3 Relationships between fissure-fluid coupling and tungsten mineralization

#### 3.1 Fissure-fillings and its compositions

Fissure-fillings species may help to reveal fissure-fluid coupling in during W mineralization. Fissure-fillings may reveal compositions of the

ore-forming fluids, and they could indicate the evolution of the ore-forming fluids. Three different occurrences of scheelite were found in the study area, and different alteration associations were recognized in the fissure-fillings. Fracture alteration mineral filling sequence respectively is iron tourmaline → microcline → Na-feldspar → biotite → siliceous → wolframite → scheelite-black tungsten → pyrrhotite → pyrite → chalcopyrite → chlorite → calcite, the filling sequence of minerals in fissure is silicate → oxide → sulphide → carbonate minerals, indicated that mineralization evolution were from high temperature to low temperature facies. Oxide phase is mainly mineralization of scheelite phase.

#### 3.2 Estimation of Fracture Permeability

Its migration path, manners and quantity of ore-forming fluid migration are controlled by the permeability of crushed rock. The permeability of cataclastic facies is a function of the cube of the fracture density and the fracture aperture (or width). Snow (1970) derived the following formula:  $K = (nd^3) / 12$ , K is the permeability (cm<sup>2</sup>); n is the crack density (strip/cm); d is the width of crushed fracture (cm).

Scheelite (table 1) is mainly hosted in biotite hornstone with higher permeability than these in biotite diorite except for near-surface (0-149.54m). Fracture density gradually reduced from surface to underground. Fracture density is usually high surrounded scheelite mineralization body. Therefore, the fissure is scheelite ore-hosting space and mineralization fluid migration channel.