

JIN Zhongguo, ZHENG Minghong, ZHOU Jiayi, HUANG Zhilong and LIU Ling, 2017. "Structure-Lithologic-Fluid" Metallogenic Coupling of the Wuzhishan Lead-Zinc Deposit in Puding, Guizhou Province. *Acta Geologica Sinica (English Edition)*, 91(supp. 1): 217-219.

"Structure-Lithologic-Fluid" Metallogenic Coupling of the Wuzhishan Lead-Zinc Deposit in Puding, Guizhou Province

JIN Zhongguo^{1,*}, ZHENG Minghong¹, ZHOU Jiayi², HUANG Zhilong² and LIU Ling¹

1 Non-Ferrous Metals and Nuclear Industry Geological Exploration Bureau of Guizhou, Guiyang 550005, China;

2 State Key Laboratory of Ore Deposit Geochemistry, Institute of Geochemistry, Chinese Academy Sciences, Guiyang 550081, China;

1 Introduction

The Wuzhishan lead-zinc ore-concentrated area in Puding is located in the east of the Sichuan, Guizhou and Yunnan lead-zinc metallogenic domain, with the Youjiang-Nanpan River metallogenic province to the south. Cambrian, Carboniferous, Permian and Triassic strata outcrop extensively in this area, with scattered distribution of Cretaceous and Sinian strata. The Upper Sinian Dengying Formation and the Lower Cambrian Qingxudong Formation are main ore-bearing strata and altered dolomite is the main host rock. Complex structure of this area mainly consists of NW-trending Ziyun-Yadou faults, NW-trending Dayuan and Wuzhishan anticline, EW-trending Nayong-Kaiyang faults, and NE-trending Anshun-Pingba faults, etc.

In recent 5 years, the first large Pb-Zn deposit(Pb-Zn reserves ≥ 1.35 Mt) has been explored in the Wuzhishan ore concentrated area, achieving geological prospecting breakthrough. Many scholars carried out research on ore-forming material source, ore deposit genesis and metallogenic regularity(Chen et al., 2015; Jin et al., 2015; Jin et al., 2016), but the research on metallogenic coupling is limited. In this paper, the "structure-lithologic-fluid" metallogenic coupling was explored on the basis of macroscopic geological investigation, combined with isotope and fluid inclusion research, aiming to deepen and enrich lead-zinc metallogenic theory in this area.

2 Typical ore deposit geological feature description--take Nayongzhi ore deposit as an example

Nayongzhi lead-zinc deposit occurred in the northeastern segment of the Wuzhishan anticline. NE- and NW-trending faults development, NE-trending faults mainly includes F₁, F₂, F₇ and etc. (ore directing and controlling faults), and NW-trending faults are the main secondary fault. Pb-Zn orebody has two types such as stratiform or stratiform-like type and vein type. The former occurs in the Qingxudong Formation dolomite at the F₇ footwall, which can be divided into I, II and III orebodies (Figure 1). The latter, steep veins, occurs in F₇ fault fracture zone. They are the products of the same fluid system in different space(Jin, et al., 2016). Mineral composition is simple, and the metal minerals were mainly sphalerite, galena, and pyrite, and gangue minerals were mainly dolomite and calcite. Ore structure contains massive, disseminated, net-vein, breccia and etc.; Ore texture contains anhedral- euhedral granular texture, fracture texture, residual texture and etc. Dolomitization and silicification are closely related to mineralization.

3 "Structure-lithologic-fluid" metallogenic coupling

3.1 Structure is necessary for lead-zinc mineralization and main ore-controlling factors

The Wuzhishan anticline structure developed. All the lead-zinc ore deposits(or spots) having been found and explored occur within the metallogenic boundary controlled by F₁ and F₂. Ore-directing and ore-controlling faults at or near the axis of the anticline control the occurrence of the main orebody. The mineralization in the F₇ fault fracture zone was strong. The thickness of ore bodies and width of faults have a positive correlation. Stratiform and stratiform-like orebodies controlled by interlayer faults distributed at the footwall

* Corresponding author. E-mail: gzkyjzg@sina.com




Fig.1 Geological map and No.0 cross section of the Nayongzhi Pb-Zn deposit

1-Lower Cambrian Jindingshan Formation; 2-First section of the Lower Cambrian Qingxudong Formation; 3-First layer of the second section of the lower Cambrian Qingxudong Formation; 4-Second layer of second section of the lower Cambrian Qingxudong Formation; 5-Third section of the lower Cambrian Qingxudong Formation; 6-Lower Carboniferous Xiangbai Formation; 7-Lower Carboniferous Dapu Formation; 8-Upper Permian Longtan Formation; 9-Lower Triassic Daye Formation; 10-Drill and number; 11-Cross section and number; 12-Thrust fault; 13-Nomal fault; 14-Steep veined ore body; 15-Stratiform ore body; 16-Dolomite; 17-Argillaceous dolomite

within the 800m range of the major fault. The closer to the F_7 fault, the higher the ore grade is, whereas lead-zinc mineralization would be diminished or vanished. Characteristics of hydrothermal mineralization and ore-controlling by structure are obvious. In short, the Wuzhishan anticline provides the entrapment structural environment for ore-forming fluid migration and ore deposition and enrichment.

3.2 Dolomite is good ore-containing Lithology

Altered coarse crystalline dolomite in the study area, as the transformation product of regional basin brine activities, is closely related to lead-zinc mineralization. Previous studies show that stronger dolomitization and higher porosity is more likely to lead to the migration of ore-forming fluid during the water-rock interaction and the lead-zinc ore precipitation, enrichment and mineralization (Han, et al., 2012; Zhen et al., 2013). Altered dolomite, spatially and temporally constrained by the metallogenic structure (Han, et al., 2014), is good ore-hosting rock. Lead-zinc mineralization was selective to certain lithology.

3.3 Ore-forming fluid properties and metallogenic dynamics background

Isotope and fluid inclusion investigation (Jin, et al., 2016) show that the ore-forming metals Pb and Zn mainly come from basement rock and ore-hosting rock. Mineralization agent C and S come from marine carbonate rock and evaporate rock at ore-hosting strata respectively. O mainly comes from deep metamorphic water. Ore-forming fluid

has the characteristics of multiple source and basin brine with low temperature and low-middle salinity. Metallogenic age in this area ranges from 200 to 230 Ma (Han, et al., 2014). The deposit formed under the geodynamic background of transition from compression to extension.

3.4 “Structure-lithologic–fluid” metallogenic coupling

Water/rock reaction between deep metamorphic water circulation driven by compressional orogeny and strata that the fluid flowed through → the ore-forming fluid formed from leaching and dissolving ore-forming metals and mineralizer migration along the fault → A massive dolomitization and gas-liquid fluid differentiation resulting from vacuum outgassing (leading to rich CO_2 fluid escaping into carbonate rocks) (Han, et al., 2014) → The opening of the structure space leads to the changes of the geochemical environment, resulting in ore-forming fluid differentiation, ore deposition and enrichment. The lead-zinc deposits in this area are the product of “Structure-lithologic–fluid” metallogenic coupling mineralization.

Acknowledgements

This research project was funded by the national key basic research development program (2014 CB440905), and key project of national natural science funds (41430315). Sincere acknowledgements should also be given!

References

- Chen Guoyong, Wang Liang, Fan Yumei and Zheng Wei . 2015. Ore-search prospect of the deep subsurface in the Wuzhishan Pb-Zn ore field, Guizhou province. *Geology and Exploration*, 51(5): 0859–0869 (in Chinese with English abstract).
- Han Runsheng, Hu Yuzhao, Wang Xuekun, Hou Baohong, Huang Zhilong, Chen Jin, Wang Feng, Wu Peng, Li Bo, Wang Hongjiang, Dong Ying and Lei Li. 2012. Mineralization modle of rich Ge—Ag—bearing Zn—Pb polymeeallic deposit concentrated district in Northeastern Yunnan,China. *Acta Geologica Sinica*, 86(2):280-294 (in Chinese with English abstract)..
- Han Runsheng, Wang Feng, Hu Yuzhao, Wang Xuekun, Ren Tiao, Qiu Wenlong and Zhong Kanghui.2014.Metallogenic tectonic dynamics and chronology constrains on theHuize-type (HZT) germanium-rich silver-zinc-lead deposits .*Geotectonica ET Metallogenia*, 38(4):758-771 (in Chinese with English abstract).
- Jin Canhai, Li Kun, Huang Lin, Zhang Yu and Shen Zhanwu. 2015. Characteristics of sulfur and lead isotope composition and metallogenic material source of the Nayongzhi Pb-Zn deposit, northwestern Guizhou province. *Journal of Mineralogy and Petrology*, 35(3): 81–88 (in Chinese with English abstract).
- Jin Zhongguo, Zhou Jiaxi Huang Zhilong, Luo Kai, Gao Jianguo, Peng Song, Wang Bing and Chen Xinglong. 2016. Ore genesis of the Nayongzhi Pb-Zn deposit, Puding city, Guizhou province, China: Evidences from S and in situ Pb isotopes. *Acta Petrologica Sinica*, 32(11): 3441–3455(in Chinese with English abstract).
- Zhen Shimin, Zhu Xinyou, Li Yongsheng Du Zezhong, Gong Xiaodong, Gong Fanying and Qi Fanyu. 2013. A tentative discussion on Mississippi valey-tpye deposits. *Mineral deposits*, 32(2): 367–379 (in Chinese with English abstract).