

LIU Qingquan, SHAO Yongjun, ZHOU Keping and LI Yongfeng, 2017. Petrogeochemistry, geochronology and its geological significance of the granite from Baishipo Ag deposit, Dabie orogen. *Acta Geologica Sinica* (English Edition), 91(supp. 1): 222-223.

## Petrogeochemistry, geochronology and its geological significance of the granite from Baishipo Ag deposit, Dabie orogen

LIU Qingquan<sup>1,2</sup>, SHAO Yongjun<sup>2\*</sup>, ZHOU KePing<sup>1</sup> and LI Yongfeng<sup>3</sup>

1 School of Resources and safety engineering, Central South University, Changsha, Hunan 410083

2 Key Laboratory of Metallogenic Prediction of Nonferrous Metals and Geological Environment Monitoring (Central South University), Ministry of Education, Changsha 410083, China

3 Henan Provincial Non-Ferrous Metals Geological and Mineral Resources Bureau, Zhengzhou, Henan 450016

### 1 Introduction

Granitoid magmas have been widely used as a natural probe for tracing the crustal evolution process (Martin et al., 2005). Mesozoic granitoids are widespread in the Dabie Orogen. Comprehensive investigations of these granitoids represent a promising opportunity to obtain deeper insight into the petrogenesis, mechanism of magma formation, geodynamic evolution and metallogeny of the associated mineral deposits. The Dabie Orogen contains tens of granitic complexes, which occur either as individual plutons with an areal extent of tens to hundreds of km<sup>2</sup> or as small stocks (generally of <1 km<sup>2</sup>) (Liu et al., 2017). The Mesozoic granitoids in the Dabie Orogen are of particular geological interest as indicators for Mesozoic lithospheric evolution.

The Baishipo Ag deposit is located in Luoshan County, Henan Province, China. In this paper, we report the geochronology and geochemistry data to define the timing of magmatism, origin of the magmas, and geodynamic setting.

### 2 Timing of magmatism

Intrusions are widely distributed in the eastern Dabie Orogen along the NW- and NNE-trending faults. The intrusive rocks are composed of granitoid plutons and minor mafic-ultramafic rocks. The Yanshannian granitoids consist of deep-seated granite batholiths, such as the Shangcheng, Xingxian and Lingshan intrusions, and other small stocks, such as those at Shapinggou, Tangjiaping, Qian'echong, Yaochong, Dayinjian, Mushan, and Xiaofan. The large range of chronometric

ages indicates that those stocks can be divided into an early stage (143–130 Ma) and a late stage (130–113 Ma) (Liu et al., 2017).

To constrain the timing of the Baishipo granite, we report the zircon U–Pb dating. Zircons from the Baishipo granite yield the weighted average U–Pb ages of 135 ± 2.0 Ma (MSWD = 1.18), we propose that magmatism were formed in the Early Cretaceous.

### 3 Genetic type and magma source

The Baishipo granite have high SiO<sub>2</sub> values, and low Mg<sup>#</sup>, Ni, and V contents, suggesting that they were originated from crustal materials. According to following evidence, We propose that the Baishipo granite is the A-type granite which have high K<sub>2</sub>O + Na<sub>2</sub>O, Ga/Al, and low CaO, Ba, Sr, and are enriched in Zr, Zn, Nb, and REE (Halen et al., 1987). It is generally accepted that A-type granites are originated from relative high-temperature magmas compared with I- and S-type granites (Patino, 1997). The calculated T<sub>Zr</sub> of the Baishipo granite samples vary from 804 to 821 °C, with an average of 810 °C, Which is similar to those of typical A-type granites (Miller et al., 2003). As discussed above, we conclude that the Baishipo granite belongs to A-type granite.

The Baishipo granite have relatively high HREE values, indicating garnet as the main residual phase (Defant et al., 1990). Additionally, the samples have high Rb/Ba ratios and low Sr contents, consistent with middle to upper continental crust (Rudnick et al., 2003). Therefore, the source region of the Baishipo granites might be relatively shallow (probably < 30 km). Experiment indicate that partial melting of calc-alkaline tonalite to granodiorite can produce peraluminous A-type

\* Corresponding author: E-mail: shaoyongjun@126.com

granitic melts on the condition of  $P = 0.4$  GPa,  $T = 950$  °C, which would lose A-type geochemical features as pressure increased up to  $> 0.8$  Gpa (Patina, 1997). Therefore, we propose that the primitive magma of the Baishipo A-type granite was originated from intermediate felsic rocks under the condition of 0.4–0.8 GPa pressures and ca. 14–28 km depths (Gao et al., 2016).

#### 4 Geodynamic setting and its implications

Zircons U–Pb age confirm that the Baishipo granite was formed in the Early Cretaceous. The Dabie Orogen is a Triassic collision zone between the North China Craton and Yangtze Block, the coesite and diamond inclusions in eclogites suggest that the continental crust subducted into mantle depths (Ames et al., 1993). Geophysical data show that the Dabie Orogen is average crust, rather than the thickened crust (Gao et al., 1998), which indicates that it has a significant thinning of the lithosphere. The Izanagi Plate (or Paleo-Pacific) started to subduct beneath the Eurasian continent in a NWW direction in the late Jurassic–Early Cretaceous period (Niu et al., 2003), which led to the transformation of tectonic regime in the Dabie Orogen. The geodynamic setting of mineralization in this period has been a NE–SW direction extension. After this period, the Izanagi Plate underwent NE-directed subduction parallel to the Eurasian continental margin, which led to large-scale continental extension (Mao et al., 2013). Therefore, the Early Cretaceous granites may have formed in a extensional setting triggered by the oblique subduction of the Izanagi Plate (Ling et al., 2009; Mao et al., 2008).

It is generally accepted that A-type granites formed in extensional environments (Alirezai et al., 2012). We infer that the Baishipo granite were formed in the post-collision extension setting combination with the regional tectonic setting and geochemical data. The thinning of the lithosphere and upwelling of asthenosphere had taken place during this extension. The direct heating effect of the asthenosphere caused the partial melting of the crust by thermal and chemical erosion. Therefore, the formation of the Baishipo A-type granite formed in the extensional setting in the Dabie orogen.

#### Acknowledgements

The study was supported by the Innovation driven project of Central South University (2015CX008). We are grateful to the ALS Mineral/ALS Chemex (Guangzhou) Co. Ltd., and Testing Research Center of Shandong Bureau, China Metallurgical Geology Bureau for the analytical work.

#### References

Alirezai, S., and Hassanzadeh, J., 2012. Geochemistry and

- zircon geochronology of the Permian A-type Hasanrobat granite, Sanandaj–Sirjan belt: a new record of the Gondwana break-up in Iran. *Lithos*, 151(11): 122–134.
- Ames, L., Tilton, G.R., and Zhou, G.Z., 1993. Timing of collision of the Sino-Korean and Yangtze Blocks: U–Pb zircon dating of coesite-bearing eclogites. *Geology*, 21(3): 339–342.
- Defant, M.J., and Drummond, M.S., 1990. Derivation of some modern arc magmas by melting of young subducted lithosphere. *Nature*, 347(6294): 662–665.
- Gao, S., Luo, T.C., Zhang, B.R., Zhang, H.F., Han, Y.W., Zhao, Z.D., and Hu, Y.k., 1998. Chemical composition of the continental crust as revealed by studies in East China. *Geochimica Et Cosmochimica Acta*, 62(11): 1959–1975.
- Gao, Y., Mao, J.W., Ye, H.S., Li, Y.F., Luo, Z.Z., and Yang, Z.Q., 2016. Petrogenesis of ore-bearing porphyry from the Tangjiaping porphyry Mo deposit, Dabie orogen: Zircon U–Pb geochronology, geochemistry and Sr–Nd–Hf isotopic constraints. *Ore Geology Reviews*, (79): 288–300.
- Halen, J.B., Currie, K.L., and Chappell, B.W., 1987. A-type granites: geochemical characteristics, discrimination and petrogenesis. *Contributions to Mineralogy and Petrology*, 95(4): 407–419.
- Ling, M.X., Wang, F.Y., Ding, X., Hu, Y.H., Zhou, L.B., Zartman, R.E., Yang, X.Y., and Sun, W.D., 2009. Cretaceous ridge subduction along the Lower Yangtze River belt, Eastern China. *Economic Geology*, 104(2): 303–321.
- Liu, Q.Q., Zhang, Y., Shao, Y.J., Zhou, K.P., Li, Y.F., Luo, Z.Z., 2017. Age, nature, and origin of early cretaceous plutons in the Dabie Mo Mineralization belt: the Xinxian example, International Geology Review, <http://dx.doi.org/10.1080/00206814.2017.1306724>
- Mao, J.W., Xie, G.Q., Bierlein, F., Qu, W.J., Du, A.D., Ye, H.S., Pirajno, F., Li, H.M., Guo, B.J., Li, Y.F., and Yang, Z.Q., 2008. Tectonic implications from Re–Os dating of Mesozoic molybdenum deposits in the East Qinling–Dabie orogenic belt. *Geochimica Et Cosmochimica Acta*, 72(18): 4607–4626.
- Mao, J.W., Cheng, Y.B., Chen, M.H., and Pirajno, F., 2013. Major types and time-space distribution of Mesozoic ore deposits in South China and their geodynamic settings. *Mineralium Deposita*, 48(3): 267–294.
- Martin, H., Smithies, R.H., Rapp, R., Moyen, J.F., and Champion, D., 2005. An overview of adakite, tonalite-trondhjemite-granodiorite (TTG), and sanukitoid: relationships and some implications for crustal evolution. *Lithos*, 79(1–2): 1–24.
- Miller, C.F., McDowell, S.M., and Mapes, R.W., 2003. Hot and cold granites? Implications of zircon saturation temperatures and preservation of inheritance. *Geology*, 31(6): 529–532.
- Niu Baogui, He Zhenjun, Song Biao, and Ren Jishun, 2003. SHRIMP dating of the Zhangjiakou volcanic series and its significance. *Geological Bulletin China*, 22(2): 140–141 (in Chinese with English abstract).
- Patino Douce, A.E., 1997. Generation of metaluminous A-type granites by low-pressure melting of calc-alkaline granitoids. *Geology*, 125(8): 743–746.
- Rudnick, R.L., and Gao, S., 2003. Composition of the continental crust. In: HOLLAND H D, TUREKIAN K K. (Eds.), *The Crust3 Treatise on Geochemistry. Elsevier-Pergamon, Oxford*, 1–64.