

MA Changqian, LIU Bin, GAO Ke and HE Zuoxiang, 2017. Petrogenesis of Triassic Mafic Complexes with MORB/OIB Affinities from the Western Garzê-Litang Ophiolitic Mélange, Central Tibetan Plateau. *Acta Geologica Sinica* (English Edition), 91(supp. 1): 19-20.

Petrogenesis of Triassic Mafic Complexes with MORB/OIB Affinities from the Western Garzê-Litang Ophiolitic Mélange, Central Tibetan Plateau

MA Changqian^{1,2*}, LIU Bin³, GAO Ke² and HE Zuoxiang²

¹ State Key Laboratory of Geological Processes and Mineral Resources, China University of Geosciences, Wuhan 430074, China

² Faculty of Earth Sciences, China University of Geosciences, Wuhan 430074, China

³ School of Geosciences, Yangtze University, Wuhan 430100, China

1 Abstract

There is a general consensus that most ophiolites formed above subduction zones (Pearce, 2003), particularly during forearc extension at subduction initiation (Shervais, 2001; Stern, 2004; Whatmam and Stern, 2011). “Supra-Subduction zone” (SSZ) ophiolites such as the well-studied Tethyan ophiolites, generally display a characteristic sequential evolution from mid-oceanic ridge basalts (MORBs) to island arc tholeiites (IATs) or bonites (BONs) (Pearce, 2003; Dilek and Furnes, 2009, 2011), which were generated in sequence from the decompression melting of asthenospheric mantle and partial melting of subduction-metasomatized depleted mantle (Stern and Bloomer, 1992; Dilek and Furnes, 2009; Whatmam and Stern, 2011). However, ophiolites with MORB and/or oceanic-island basalt (OIB) affinities are rare, and their origin and tectonic nature are poorly understood (Boedo et al., 2013; Saccani et al., 2013). It is interesting that the composition of these ophiolites from the central Tibetan Plateau (CTP) is dominated by MORBs and minor OIBs and a distinct lack of IATs and BONs, which is inconsistent with most ophiolites worldwide (Robinson and Zhou, 2008; Zhang et al., 2008). But the generation and tectonic nature of these ophiolites are still controversial.

In this study, we present new geochronological, mineralogical and Sr-Nd isotopic data for the Chayong and Xiewu mafic complexes in the western Garzê-Litang suture zone (GLS), a typical Paleo-Tethyan suture crossing the CTP (Fig. 1). The Triassic ophiolite in the western GLS has been described by Li et al. (2009), who found

that it mainly consists of gabbros, diabases, pillow basalts and a few metamorphic peridotites. The ophiolite has been tectonically dismembered and crops out in Triassic clastic rocks and limestones as tectonic blocks. The Chayong and Xiewu mafic complexes are generally regarded as important fragments of the Triassic ophiolites (e.g., Jin, 2006; Li et al., 2009). Zircon LA-ICP-MS U-Pb ages of 234 ± 3 Ma and 236 ± 2 Ma can be interpreted as formation times of the Chayong and Xiewu mafic complexes, respectively. The basalts and gabbros of the Chayong complex exhibit enriched MORB (E-MORB) compositional affinities except for a weak depletion of Nb, Ta and Ti relative to the primitive mantle, whereas the basalts and gabbros of the Xiewu complex display distinct E-MORB and OIB affinities. The geochemical features suggest a probable fractionation of olivine \pm clinopyroxene \pm plagioclase as well as insignificant crustal contamination.

The geochemical and Sr-Nd isotopic data reveal that the Chayong mafic rocks may have been derived from depleted MORB-type mantle metasomatized by crustal components and Xiewu mafic rocks from enriched lithospheric mantle metasomatized by OIB-like components. The ratios of Zn/Fet, La/Yb and Sm/Yb indicate that these mafic melts were produced by the partial melting of garnet + minor spinel-bearing peridotite or spinel \pm minor garnet-bearing peridotite. We propose that back-arc basin spreading associated with OIB/seamount recycling had occurred in the western GLS at least since the Middle Triassic times, and the decompression melting of the depleted MORB-type asthenosphere mantle and partial melting of sub-continental lithosphere were metasomatized by plume-related melts, such as OIBs, which led to the generation of the Chayong and Xiewu mafic melts.

* Corresponding author. E-mail: cqma@cug.edu.cn

Acknowledgements

This study was financially supported by the National Nature Science Foundation of China (Grant 41272079).

References

- Boedo, F.L., Vujovich, G.I., Kay, S.M., Ariza, J.P., and Pérez Luján, S.B., 2013. The E-MORB like geochemical features of the early Paleozoic mafic-ultramafic belt of the Cuyania terrane, western Argentina. *Journal of South American Earth Sciences*, 48: 73–84.
- Dilek, Y., and Furnes, H., 2009. Structure and geochemistry of Tethyan ophiolites and their petrogenesis in subduction rollback systems. *Lithos*, 113: 1–20.
- Dilek, Y., and Furnes, H., 2011. Ophiolite genesis and global tectonics: geochemical and tectonic fingerprinting of ancient oceanic lithosphere. *Geological Society of America Bulletin*, 123: 387–411.
- Jin, G.S., 2006. Geochronology and Geochemistry Characters on Some Magmatic Rocks in the West of Xijir Ulan-Jinshajiang Suture Zone. *Chinese Academy of Geological Sciences*, 1–78 (in Chinese with English Abstract).
- Li, S.P., Chen, Y.S., Li, W.F., Wang, Q.Y., and Qi, S.S., 2009. Geochemistry of the Chayong ophiolite in North Qiangtang Basin, Qinghai-Tibet Plateau: implications for tectonics. *Geoscience*, 23: 1030–1038 (in Chinese with English Abstract).
- Pearce, J.A., 2003. Supra-subduction zone ophiolites: the search for modern analogues. In: Dilek, Y., Newcomb, S. (eds.), *Ophiolite Concept and the Evolution of Geological Thought*. Geological Society of America Special Paper 373. *Geological Society of America, Boulder, CO*, 269–293.
- Robinson, P.T., and Zhou, M., 2008. The origin and tectonic setting of ophiolites in China. *Journal of Asian Earth Sciences*, 32: 301–307.
- Saccani, E., Azimzadeh, Z., Dilek, Y., and Jahangiri, A., 2013. Geochronology and petrology of the early carboniferous Misho mafic complex (NW Iran), and implications for the melt evolution of paleo-Tethyan rifting in western Cimmeria. *Lithos*, 162–163: 264–278.
- Shervais, J.W., 2001. Birth, death, and resurrection: the life cycle of suprasubduction zone ophiolites. *Geochemistry, Geophysics, Geosystems*, 2.
- Stern, R., 2004. Subduction initiation: spontaneous and induced. *Earth and Planetary Science Letters*, 226: 275–292.
- Stern, R.J., and Bloomer, S.H., 1992. Subduction zone infancy: examples from the Eocene Izu-Bonin-Mariana and Jurassic California arcs. *Geological Society of America Bulletin*, 104: 1621–1636.
- Whattam, S.A., and Stern, R.J., 2011. The ‘subduction initiation rule’: a key for linking ophiolites, intra-oceanic forearcs, and subduction initiation. *Contributions to Mineralogy and Petrology*, 162: 1031–1045.
- Zhang, Q., Wang, C.Y., Liu, D., Jian, P., Qian, Q., Zhou, G., and Robinson, P.T., 2008. A brief review of ophiolites in China. *Journal of Asian Earth Sciences*, 32: 308–324.




Fig. 1. Location of the Tibetan Plateau in the Tethyan realm (a) and simplified geological map of the TP (b).