四川乐山震旦系灯影组火山碎屑岩锆石 LA-ICP-MS U-Pb 定年及盆地裂陷演化讨论

资金平¹⁾, 贾东¹⁾, 魏国齐²⁾, 杨振宇³⁾, 张勇¹⁾, 胡晶¹⁾, 沈淑鑫¹⁾

1) 南京大学内生金属矿床成矿机制研究国家重点实验室, 南京, 210023;

2) 中国石油勘探开发研究院廊坊分院, 河北廊坊, 065007; 3) 首都师范大学, 北京, 100048

内容提要:四川盆地中部晚震旦(埃迪卡拉)世—早寒武世克拉通内裂陷与安岳震旦(埃迪卡拉)系—寒武系特大气田的形成关系密切,年代学研究对认识克拉通内裂陷构造意义重大。对四川盆地乐山先锋剖面灯影组三段火山碎屑岩进行锆石 LA-ICP-MS U-Pb 定年,研究结果表明四川盆地灯影组三段火山碎屑岩的锆石 U-Pb 年龄为(539.6 ± 1.4) Ma,年代学结果表明四川盆地灯影组三段与灯影组四段属于寒武系。基于灯影组底界(551.1 Ma)、含火山碎屑岩的灯影组三段(539.6 Ma)、含小壳化石的麦地坪组(535.2 Ma)和下寒武统 Ni—Mo 多金属硫化物富集层(521.0 Ma)四个等时面构建了克拉通内裂陷演化的时间框架。在此基础上认为四川盆地中部晚震旦(埃迪卡拉)世—早寒武世隆坳构造的形成主要受边界断裂控制,断层活动使裂陷内水体加深,少量灯影组四段含硅质白云岩沉积后,麦地坪组与筇竹寺组开始沉积,麦地坪组与灯影组四段的界线在裂陷内与裂陷两侧并不等时,裂陷内麦地坪组、筇竹寺组与裂陷两侧灯影组四段为同期异相沉积。

关键词:四川盆地;寒武纪;震旦(埃迪卡拉)纪;灯影组;锆石定年;克拉通内裂陷

四川盆地油气储量丰富,开采潜力巨大(王世谦等, 2005; 李世臻等, 2013; 何洪春等, 2015; 邹才能等, 2015)。2011年7月,川中高石1井在寒武系龙王庙组与震旦(埃迪卡拉)系灯影组取得重大勘探突破(杜金虎等, 2014)。位于四川盆地中部呈近南北向展布的隆坳构造格局是形成区域有利勘探条件的关键因素。钻井与地震剖面资料显示,坳陷内震旦系灯影组三段与灯影组四段厚度远小于坳陷两侧,而麦地坪组与筇竹寺组厚度远大于坳陷两侧。目前对该隆坳构造格局的成因尚存在多种解释方案,汪泽成等(2014)认为震旦纪末期桐湾运动Ⅱ幕区域逆冲—褶皱隆升、剥蚀作用以及早寒武世拉张背景是形成隆坳格局的关键。李忠权等(2015)认为早寒武世地壳差异抬升运动,河流下切侵蚀和溶蚀作用是坳陷内灯影组三段与灯影组四段厚度薄的关键,随后坳陷内沉积充填了厚层的下寒武统麦地坪组和筇竹寺组。魏国齐等(2015)提出隆坳构造为震旦纪—早寒武世克拉通内裂陷,认为灯影组三段、灯影组四段沉积期,裂陷内呈现欠补偿沉积环境,仅沉积了薄层灯

三段碎屑岩与灯影组四段含硅质白云岩。

明确隆坳构造格局的成因机制对研究四川盆地深部油气勘探方向具有十分重要的意义。普遍含火山碎屑岩的灯影组三段为四川盆地灯影组对比的重要标志层(邓胜徽等, 2015),亦是隆坳构造格局的关键层位。本文对四川乐山先锋剖面灯影组三段火山碎屑岩进行锆石 LA-ICP-MS U-Pb 定年,以期对隆坳构造格局增添新的认识。

1 区域地质背景

震旦纪至早寒武世,华南扬子地块具有自西北向东南水体总体加深的趋势(Zhu Maoyan et al., 2003; Steiner et al., 2007)。古地理可依次划分为内部台地带、过渡带和斜坡至深水盆地带,界线沿北东—南西展布。沉积环境差异引起区域沉积发生相变,晚震旦世浅水台地相区以沉积碳酸盐岩为主,而在水体加深的过渡带和斜坡至深水盆地带则普遍沉积碳质硅质岩。进入寒武纪,扬子地块普遍沉积了一套黑色页岩(陈南生等, 1982),上扬子地区该套

注:本文为国家科技重大专项(编号:2016ZX05003-001、2017ZX05008-001)和国家自然科学基金资助项目(编号:41230208)的成果。

收稿日期:2016-12-19;改回日期:2017-05-10;责任编辑:刘志强。10.16509/j.georeview.2017.04.014

作者简介:资金平,男,1991年生。硕士研究生,构造地质学专业。zijinping@qq.com。通讯作者:贾东,男,1960年生。博士,教授,主要从事构造地质学研究。Email: djia@nju.edu.cn。

图1 四川盆地震旦纪(埃迪卡拉纪)灯影组二段沉积期末和灯影组四段沉积期末裂陷分布范围图(据魏国齐等,2015 修改)

Fig. 1 The rift areas at the end deposition time of the Second and the Fourth Member of the Sinian (Ediacaran) Dengying Formation((modified after Wei Guoqi et al. , 2015&)

黑色页岩底部或之下为一套富磷地层(陈志明等,1987;牟南等,2005;毛铁等,2015)。四川盆地处于扬子地块西部的内部台地带,晚震旦世一早寒武世水体较浅,自下而上地层沉积了以白云岩为主的灯影组、以含磷白云岩为主的麦地坪组、以黑色碳质页岩为特征的筇竹寺组、下部碎屑岩段上部碳酸盐岩段的沧浪铺组以及龙王庙组海相碳酸盐岩。

扬子地块震旦系上统与下统界线位于陡山沱组内部(Zhu Maoyan et al. , 2007; 刘鹏举等, 2012; 全国地层委员会《中国地层表》编委会, 2014)。邓胜徽等(2015)将四川盆地震旦系上统灯影组自下而上划分为四个岩性段:以贫菌藻类白云岩为特征的灯一段;以下部富含菌藻类、上部贫菌藻类白云岩为特征的灯二段;岩性为碎屑岩,常夹火山碎屑岩层的灯三段;以硅质条带为特征的灯四段。罗贝维等(2013a,b)研究指出,四川盆地灯影组沉积期,海水温度介于 $15 \sim 30^{\circ}\text{C}$,灯一段沉积期海平面逐渐上升;灯二段沉积期海平面明显降低且整体处于动荡的海退环境;灯三段沉积期,发生了一次范围大、时间短的海侵事件;灯四段沉积期,海平面逐渐降低,

并在后期存在持续震荡的过程。

震旦纪末至早寒武世的桐湾运动与川中隆拗构造格局的形成关系紧密(汪泽成等,2014;武赛军等,2016)。三幕桐湾运动引起的区域抬升与剥蚀分别形成了四川盆地灯影组三段与二段之间、下寒武统麦地坪组与灯影组四段之间以及筇竹寺组与麦地坪组之间的普遍假整合,也使灯影组二段与四段发育大量溶蚀孔洞,成为优质储层(汪泽成等,2014)。而在区域上,同一幕运动在整体幕式隆升的背景下,局部地区存在差异升降(武赛军等,2016)。

川中隆拗构造格局呈南北向展布,东部边界陡,西部边界缓(图1)。拗陷内震旦系一下寒武统沉积存在以下三个特点:①通过钻井标定后的地震解释预测拗陷内灯二段厚度略小于拗陷东西两侧灯二段厚度,位于拗陷东侧的高石1井灯二段厚度约为486.8 m;②灯三段碎屑岩、灯四段白云岩厚度薄,以高石17井为例,灯三段厚度约2.0 m,灯四段厚度约15.0 m;拗陷东侧高石1井,灯四段厚262.5 m;③拗陷内麦地坪组和筇竹寺组黑色页岩和浅灰色泥岩累计厚度介于500~700 m,大于东西两侧约300~500 m(魏国齐等,2015)。隆拗构造的成因目前尚有不同观点(汪泽成等,2014;李忠权等,2015;魏国齐等,2015),关键层位的年代学约束至关重要。

谷志东等(2015)获得四川盆地威远地区威117井基底花岗岩锆石 SHRIMP U-Pb 年龄为 (794 ± 11) Ma,指示了区域基底结晶时间。Condon 等(2005)获得湖北宜昌峡东震旦系陡山沱组底部与顶部凝灰岩的高精度锆石 U-Pb 年龄分别为 (635.2 ± 0.6) Ma 和 (551.1 ± 0.7) Ma,以上两个年龄约束了扬子地块陡山沱组与灯影组的起始沉积时间。麦地坪组为四川盆地小壳化石的首现层位(陈孟羲,1982),朱日祥等(2009)获得云南梅树村剖面小壳化石首现层位附近凝灰岩(即第5层)年龄为 (535.2 ± 1.7) Ma,该年龄亦可大致约束四川盆地麦地坪组的沉积时间。Compston 等(2008)获得四川乐山范店剖面筇竹寺组底部凝灰岩锆石 SHRIMP U-Pb 年龄 (526.5 ± 1.1) Ma,该年龄约束了区域筇竹寺组的沉积时间。华南扬子地块下寒武统具一 Ni、Mo、PGE 和 Au 等多金属硫化物富集层,延伸长度可达1600 km,Zhu Maoyan 等(2003)提出将其作为华南下寒武统划分与对比的地球化学标志层。在四川盆地,该富集层在四川盆地筇竹寺组(或郭家坝组)底

白云岩
dolomite	泥岩
mudstone	炭质泥岩
carbon mudstone	粉砂质泥岩
silty mudstone	石英砂岩
quartz sandstone
含磷白云岩
phosphoric dolomite	砂质白云岩
sandy dolomite	硅质白云岩
siliceous dolomite	溶蚀孔洞
solution pore	葡萄状构造白云岩
grape-like dolomite
含藻白云岩
algae dolomite	硅质条带
siliceous band	花岗岩
granite	页岩
shale	火山碎屑岩
volcanic ash beds

图 2 四川乐山先锋剖面图及部分野外露头照片

Fig. 2 Geological sketch map of the Xianfeng Section in Leshan, Sichuan and some photo of sampling layers

- (a) 四川乐山先锋剖面图;(b) 采样位置示意图。左图为采样层露头,右图为采样层露头素描图;(c) 样品 XF22-S5 采样露头照片;(d) 样品 XF22-S6 采样露头照片;(e) 样品 XF22-S5 显微镜下照片;矿物颗粒极细,能识别石英(Qtz)、绿帘石(Ep)等矿物。(f) 样品 XF22-S6 显微镜下照片。火山碎屑结构,岩屑与晶屑含量超过 80%,矿物含角闪石(Hbl)、石英(Qtz)等
- (a) Geological sketch map of the Xianfeng Section in Leshan, Sichuan province; (b) Photo showing the sampling site and the sketch map is in the right; (c) Photo showing the sampling layer of the Sample XF22-S5; (d) Photo showing the sampling layer of the Sample XF22-S6; (e) Photomicrograph of the Sample XF22-S5, the grains are small, the quartz and epidote are recognized; (f) Photomicrograph of the Sample XF22-S6, Hornblende, quartz and pyroclastic texture are recognized, debris weighted more than 80 percent

部亦有沉积记录(Guo Qingjun et al., 2007)。Xu Lingang 等(2011)获得该 Ni—Mo 多金属硫化物富集层年龄为 (521 ± 5) Ma。

先锋剖面位于四川省乐山市峨边县北东向约 20 km 的先锋村与老鸦村之间(图 1),是四川盆地灯影组的典型剖面,位于坳陷西侧。前人对先锋剖面进行了丰富的岩石地层学、同位素化学地层学和同位素年代学研究并应用于古环境恢复、四川盆地及周缘地区地层对比和古地理重建(罗贝维等, 2013b; 邓胜徽等, 2015; 刘静江等, 2015)。研究区陡山沱组(Z_1-2d)直接覆盖于前震旦花岗岩基底之上,灯影组与陡山沱组碎屑岩整合接触,下寒武统麦地坪组(ϵ_1m)与灯影组接触关系为平行不整合(图 2a)。剖面灯二段与灯三段之间见有起伏不平的不整合面(图 2b)。我们于先锋剖面靠近灯三段底部位置发现两层火山碎屑岩,前者厚度约 3 cm,后者厚度不足 1 cm,层位相距约 6 cm(图 2b、c、d)。分别取样标记为 XF22-S5 与 XF22-S6 以进行锆石 U-Pb 年代学研究,采样过程中严格避免围岩污染。

2 分析方法与实验结果

用常规的重液和磁选方法分选样品中的锆石,锆石挑选完成后在双目镜下将挑选的锆石粘在双面胶上,用环氧树脂与三乙醇胺(7:1)的混合胶固结,在 60℃ 恒温状态下固化;并用 7000 目砂纸将锆石打磨直至大部分锆石暴露出内部晶面;对待测锆石做透射光、反射光和阴极发光(CL)图像分析(图 3),以确定锆石形态,揭示锆石内部结构并帮助选定最佳的锆石分析位置。锆石定年分析在南京大学内生金属矿床成矿机制研究国家重点实验室完成,仪器为美国 Agilent 公司的 7500a 型四极杆电感耦合等离子体质谱仪(ICP-MS)。激光剥蚀系统采用美国 New Wave 公司的 UP-213 Laser Ablation System 固体激光器,激光束斑直径为 24 μm 。测试分析标样为 GJ-1 (608.5 ± 1.5 Ma) 和 Mud Tank (732 ± 5

Ma)(Jackson et al., 2004)。单点分析的同位素比值及年龄误差为 1σ ,加权平均年龄误差为 95% 置信度。运用 GLITTER 4.4.1 软件包计算各测点的 U-Pb 年龄,测试原理、数据处理和铅校正方法详见 Griffin 等(2004)。数据处理采用 Isoplot/Ex v. 4.15 软件(Ludwig, 2008)。

挑选样品 XF22-S5 中 31 颗与样品 XF22-S6 中 9 颗晶形完整、岩浆生长环带清晰的锆石进行 U-Pb 同位素年龄分析。所分析的锆石 Pb 含量介于 32 ~ 343 $\mu\text{g/g}$, U 含量介于 51 ~ 1239 $\mu\text{g/g}$, Th 含量介于 62 ~ 1110 $\mu\text{g/g}$, Th/U 比值介于 0.59 ~ 2.79, 平均值 1.28。

40 个分析点共获得 38 个谐和年龄(图 4),有

图 3 先锋剖面灯三段火山碎屑岩层的代表性锆石阴极发光(CL)图像

Fig. 3 CL image of typical zircons from volcanic ash beds in the Third Member of the Dengying Formation in the Xianfeng Section

图中白色圆圈表示样品 XF22-S5 锆石定年分析位置,黑色圆圈表示样品 XF22-S6 锆石定年分析位置

The white circles mean analysis locations of zircons from the Sample XF22-S5, the black circles mean analysis locations of zircons from the Sample XF22-S6

图4 先锋剖面灯三段火山碎屑岩层的锆石 LA-ICP-MS U-Pb 年龄分析结果

Fig. 4 LA-ICP-MS U-Pb dating result of zircons from volcanic ash beds in the Third Member of the Dengying Formation in the Xianfeng Section

两个不谐和年龄,这可能是 Pb 丢失的结果,大部分锆石颗粒形成后 U-Pb 同位素体系保持封闭。样品 XF22-S5 中 17 个最年轻年龄与样品 XF22-S6 中 4 个最年轻年龄一致,本文对其进行合并分析(表 1)。21 个年龄集中分析点的谐和年龄为 (539.6 ± 1.4) Ma(图 4),加权平均年龄为 (539.4 ± 3.0) Ma(图 5)。根据锆石的形态、Th/U 比值和 CL 图像特征,我们将这 21 个分析点的 U-Pb 谐和年龄 (539.6 ± 1.4) Ma 解释为灯三段火山碎屑岩层的锆石结晶年龄。

XF22-S6-5 具有核幔构造,边部年龄 (539 ± 5) Ma,揭示了锆石结晶年龄,内部年龄为 (805 ± 7) Ma。除 21 个集中年龄外,17 个谐和年龄分布在 700 ~ 900 Ma 之间,与扬子地块中广泛存在的 0.75 ~ 0.83 Ga 新元古代岩浆岩年龄基本一致(Li ZX et al., 2003),推测为捕获锆石。

3 讨论

3.1 四川盆地灯三段火山碎屑岩层年龄及意义

本文获得四川盆地灯三段火山碎屑岩层的锆石 LA-ICP-MS U-Pb 年龄为 (539.6 ± 1.4) Ma,接近全球寒武纪底界年龄 (541 ± 1) Ma,指示了采样层位位于震旦纪—寒武纪界线附近。近期,Chen Daizhao 等(2015)获得贵州铜仁坝黄剖面、湖南张家界柑子坪剖面留茶坡组凝灰岩夹层的锆石 SHRIMP 年龄分别为 (542.6 ± 3.7) Ma 和 (542.1 ± 5.0) Ma,与本

图5 先锋剖面灯三段火山碎屑岩层的锆石 $^{206}\text{Pb}/^{238}\text{U}$ 年龄加权平均计算结果

Fig. 5 Weighted mean $^{206}\text{Pb}/^{238}\text{U}$ dating result of zircons from volcanic ash beds in the Third Member of the Dengying Formation in the Xianfeng Section

实心符号表示样品 XF22-S5,
空心符号表示样品 XF22-S6

The filled symbol means ages from the Sample XF22-S5,
and the other means ages from the Sample XF22-S6

文所获年龄在误差范围内是一致的,指示着四川盆地灯三段的火山碎屑岩层在扬子地块分布广泛。

年代学研究成果表明灯三段火山碎屑岩层的锆石 U-Pb 年龄将四川盆地震旦系—寒武系界线约束在灯三段附近。灯三段层位低于小壳化石首现层位麦地坪组约 200 m,与寒武系全球界线层型纽芬兰

剖面相类似。全球前寒武系—寒武系界线定义于纽芬兰剖面遗迹化石 *T. pedum* 首现层位 (Brasier et al., 1994), 该位置距小壳化石首现点亦有约 150 m 的寒武纪地层。

3.2 对四川盆地震旦纪—早寒武世

克拉通内裂陷的新认识

前人对四川盆地震旦纪—早寒武世隆拗构造格局形成机制存在的分歧在于解释拗陷内与拗陷两侧灯三段与灯四段、麦地坪组与筇竹寺组的分布特征。李忠权等 (2015) 将隆拗构造定义为拉张侵蚀槽; 魏国齐等 (2015) 提出隆拗构造为震旦纪—早寒武世克拉通内裂陷, 认为灯三段和灯四段沉积期, 裂陷内水体较深, 欠补偿沉积环境使得裂陷内仅沉积了少量灯三段与灯四段, 并绘制了灯二段沉积期末与灯四段沉积期末的裂陷范围 (图 1)。杜金虎等 (2016) 基于深层地震资料发现, 裂陷分布受断裂控制, 裂陷内部及两侧发育 NW 向为主的张性断层, 其中灯影组三段底界断距最大在 400 ~ 500 m, 向上至

沧浪铺组断距减小, 多数断层消失在龙王庙组。以上资料表明, 断层的主要活动时间在灯影组沉积期至沧浪铺组沉积期。

基于本文年代学结果我们将研究区震旦系—寒武系界线置于灯影组三段, 并以灯影组底界 (551.1 ± 0.7 Ma) (Condon et al., 2005)、含火山碎屑岩的灯三段 (539.6 ± 1.4 Ma) (本文)、小壳化石首现层位麦地坪组 (535.2 ± 1.7 Ma) (朱日祥等, 2009) 和扬子地块下寒武统 Ni—Mo 多金属硫化物富集层 (521 ± 5 Ma) (Xu Lingang et al., 2011) 构建了四个等时面以对裂陷进行系统的分析 (图 6)。我们认为受边界断裂活动影响, 裂陷内水体加深, 同期异相沉积是裂陷内与裂陷两侧下寒武统分布差异的关键所在。桐湾运动 I 幕结束后, 研究区继续接受灯影组沉积, 在边界断裂活动的影响下, 裂陷内沉积了约 17 m 厚灯三段与灯四段后即开始沉积深水陆棚相的麦地坪组泥页岩, 此时裂陷两侧台缘地区尚还在沉积灯四段含硅质白云岩, 我们推测含小壳化石的

表 1 四川乐山先锋剖面灯影组三段火山碎屑岩层的岩浆锆石 U-Pb 同位素年龄分析结果

Table 1 U-Pb isotopic data of the zircons from the volcanoclastic beds of the Third Member, Dengying Formation, in the Xianfeng section, Leshan, Sichuan

测点号	元素含量 ($\times 10^{-6}$)			Th/U	同位素比值						同位素年龄 (Ma)			谐和度 (%)
	Pb	Th	U		$\frac{n(^{207}\text{Pb})}{n(^{206}\text{Pb})}$		$\frac{n(^{207}\text{Pb})}{n(^{235}\text{U})}$		$\frac{n(^{206}\text{Pb})}{n(^{238}\text{U})}$		$\frac{n(^{207}\text{Pb})}{n(^{206}\text{Pb})}$	$\frac{n(^{207}\text{Pb})}{n(^{235}\text{U})}$	$\frac{n(^{206}\text{Pb})}{n(^{238}\text{U})}$	
					测值	$\pm\%$	测值	$\pm\%$	测值	$\pm\%$	测值 $\pm 1\sigma$	测值 $\pm 1\sigma$	测值 $\pm 1\sigma$	
XF22-S5-1	338	845	726	1.16	0.0592	2.0	0.7132	2.1	0.0874	1.4	575 $\pm$ 44	547 $\pm$ 9	540 $\pm$ 7	101
XF22-S5-2	104	302	236	1.28	0.0581	5.0	0.7025	4.9	0.0879	2.2	532 $\pm$ 113	540 $\pm$ 20	543 $\pm$ 12	99
XF22-S5-3	184	368	376	0.98	0.0584	2.4	0.7076	2.4	0.0880	1.4	545 $\pm$ 53	543 $\pm$ 10	544 $\pm$ 7	100
XF22-S5-4	202	328	416	0.79	0.0582	1.9	0.6977	1.9	0.0870	1.3	538 $\pm$ 43	537 $\pm$ 8	538 $\pm$ 7	100
XF22-S5-5	285	563	658	0.86	0.0584	1.9	0.7043	2.0	0.0875	1.4	543 $\pm$ 47	541 $\pm$ 8	541 $\pm$ 7	100
XF22-S5-6	151	376	293	1.28	0.0581	2.7	0.6935	2.6	0.0866	1.4	534 $\pm$ 60	535 $\pm$ 11	535 $\pm$ 7	100
XF22-S5-7	80	133	165	0.81	0.0595	2.2	0.7188	2.2	0.0876	1.4	586 $\pm$ 50	550 $\pm$ 9	542 $\pm$ 7	101
XF22-S5-8	321	636	767	0.83	0.0583	2.7	0.7008	2.8	0.0872	1.6	540 $\pm$ 61	539 $\pm$ 12	539 $\pm$ 8	100
XF22-S5-9	131	199	277	0.72	0.0583	2.3	0.7026	2.3	0.0874	1.4	541 $\pm$ 51	540 $\pm$ 10	540 $\pm$ 7	100
XF22-S5-10	32	64	63	1.01	0.0582	5.5	0.6977	5.3	0.0870	2.1	538 $\pm$ 125	537 $\pm$ 22	537 $\pm$ 11	100
XF22-S5-11	95	156	224	0.70	0.0602	6.4	0.7171	6.1	0.0864	2.5	611 $\pm$ 142	549 $\pm$ 26	534 $\pm$ 13	103
XF22-S5-12	249	637	576	1.11	0.0584	2.1	0.7058	2.2	0.0877	1.5	543 $\pm$ 47	542 $\pm$ 9	542 $\pm$ 8	100
XF22-S5-13	170	297	419	0.71	0.0583	2.4	0.7056	2.5	0.0878	1.6	541 $\pm$ 55	542 $\pm$ 11	543 $\pm$ 8	100
XF22-S5-14	250	647	600	1.08	0.0575	2.5	0.6963	2.6	0.0878	1.6	512 $\pm$ 57	537 $\pm$ 11	542 $\pm$ 8	99
XF22-S5-15	65	143	129	1.11	0.0576	3.8	0.6890	3.7	0.0868	1.7	515 $\pm$ 86	532 $\pm$ 15	536 $\pm$ 9	99
XF22-S5-16	270	520	621	0.84	0.0588	5.0	0.7080	4.8	0.0873	2.2	561 $\pm$ 111	544 $\pm$ 20	540 $\pm$ 11	101
XF22-S5-17	328	1017	732	1.39	0.0568	2.9	0.6895	2.9	0.0881	1.7	483 $\pm$ 65	533 $\pm$ 12	544 $\pm$ 9	98
XF22-S6-1	343	961	773	1.24	0.0591	1.8	0.7085	1.5	0.0870	0.9	570 $\pm$ 39	544 $\pm$ 6	538 $\pm$ 4	101
XF22-S6-2	129	237	335	0.71	0.0592	3.1	0.7111	2.9	0.0871	1.3	575 $\pm$ 68	545 $\pm$ 12	539 $\pm$ 6	101
XF22-S6-3	319	834	737	1.13	0.0583	1.8	0.7010	1.5	0.0872	0.9	540 $\pm$ 40	539 $\pm$ 6	539 $\pm$ 5	100
XF22-S6-4	145	417	334	1.25	0.0581	2.1	0.6947	1.9	0.0868	1.0	532 $\pm$ 48	536 $\pm$ 8	537 $\pm$ 5	100

图6 过四川盆地中部克拉通内裂陷连井剖面图 (龙王庙组底界拉平, 剖面位置见图1, 修改自魏国齐等, 2015)

Fig. 6 Well-tie Section across the intracratonic rift in the Middle Sichuan Basin (the bottom of the Longwangmiao Formation is leveled, the position of this section is shown in Fig. 1, modified after Weiguoqi et al., 2015 &)

麦地坪组沉积等时面在裂陷内位于筇竹寺组内部,表征裂陷内麦地坪组、筇竹寺组与裂陷两侧灯四段为同期异相沉积(图6)。Ni—Mo 硫化物富集层位于四川盆地筇竹寺组(或郭家坝组)底部,在裂陷内该界线位于筇竹寺组中部。等时面之间的地层厚度变化表现出同构造沉积特征。

4 结论

(1)对来自四川乐山先锋剖面灯影组三段的火山碎屑岩进行锆石 LA-ICP-MS U-Pb 定年,获得谐和年龄为 (539.6 ± 1.4) Ma,年代学结果表明四川盆地灯影组三段和灯影组四段属于寒武系。

(2)基于灯影组底界、含火山碎屑岩的灯影组三段、含小壳化石的麦地坪组和下寒武统 Ni—Mo 多金属硫化物富集层四个等时面构建了克拉通内裂陷的时间框架。在此基础上认为四川盆地中部晚震旦(埃迪卡拉)世一早寒武世隆坳构造的形成主要受边界断裂控制,断层活动使裂陷内水体加深,少量灯影组四段含硅质白云岩沉积后,麦地坪组与筇竹寺组开始沉积,麦地坪组与灯影组四段的界线在裂陷内与裂陷两侧并不等时,裂陷内麦地坪组、筇竹寺组与裂陷两侧灯影组四段为同期异相沉积。

致谢:衷心感谢西南油气田杨西南工程师在野外工作中给予的指导,感谢南京大学王孝磊教授、武兵工程师在锆石测年实验中给予的热情帮助。

参 考 文 献 / References

(The literature whose publishing year followed by a “&” is in Chinese with English abstract; the literature whose publishing year followed by a “#” is in Chinese without English abstract)

陈孟莪. 1982. 四川峨眉麦地坪剖面震旦系—寒武系界线的新认识及有关化石群的记述. 地质科学, (3): 253 ~ 264.

陈南生, 杨秀珍, 刘德汉, 肖学军, 范德廉, 王连芳. 1982. 我国南方下寒武统黑色岩系及其中的层状矿物. 矿床地质, (2): 39 ~ 51.

陈志明, 陈其英. 1987. 扬子地台早寒武世梅树村早期的古地理及其磷块岩展布特征. 地质科学, (3): 246 ~ 257.

邓胜徽, 樊茹, 李鑫, 张师本, 张宝民, 卢远征. 2015. 四川盆地及周缘地区震旦(埃迪卡拉)系划分与对比. 地层学杂志, 39(3): 239 ~ 254.

杜金虎, 汪泽成, 邹才能, 徐春春, 沈平, 张宝民, 姜华, 黄士鹏. 2016. 上扬子克拉通内裂陷的发现及对安岳特大型气田形成的控制作用. 石油学报, 37(1): 1 ~ 16.

杜金虎, 邹才能, 徐春春, 何海清, 沈平, 杨跃明, 李亚林, 魏国齐, 汪泽成, 杨雨. 2014. 川中古隆起龙王庙组特大型气田战略发现与理论技术创新. 石油勘探与开发, 41(3): 268 ~ 277.

谷志东, 翟秀芬, 袁苗. 2015. 四川盆地基底花岗岩岩锆石 SHRIMP U-Pb 定年、地球化学特征及其构造环境. 地质学报, 89(增): 296 ~ 296.

何洪春, 屈晓艳. 2015. 四川盆地页岩气资源勘探技术现状与展望. 地质论评, 61(增): 139 ~ 141.

罗贝维, 魏国齐, 杨威, 董才源. 2013a. 四川盆地灯影期海平面变化精细刻画研究. 地质论评, 59(增): 221 ~ 222.

罗贝维, 魏国齐, 杨威, 董才源. 2013b. 四川盆地晚震旦世古海洋环境恢复及地质意义. 中国地质, 40(4): 1099 ~ 1111.

刘静江, 李伟, 张宝民, 周慧, 袁晓红, 单秀琴, 张静, 邓胜徽, 谷志东, 樊茹, 王拥军, 李鑫. 2015. 上扬子地区震旦纪沉积古地理. 古地理学报, 17(6): 735 ~ 753.

刘鹏举, 尹崇玉, 陈寿铭, 李猛, 高林志, 唐烽. 2012. 华南峡东埃迪卡拉(震旦)纪年代地层划分初探. 地质学报, 86(6): 849 ~ 866.

李世臻, 周新桂, 杨海锐. 2013. 中国油气资源潜力和战略调查方向. 地质论评, 59(增): 737 ~ 738.

李忠权, 刘记, 李应, 杭文艳, 洪海涛, 应丹琳, 陈晓, 刘冉, 段新国, 彭戟. 2015. 四川盆地震旦系威远—安岳拉张侵蚀槽特征及形成演化. 石油勘探与开发, 42(1): 26 ~ 33.

牟南, 吴朝东. 2005. 上扬子地区震旦—寒武纪磷块岩岩石学特征及成因分析. 北京大学学报(自然科学版), 41(4): 551 ~ 562.

毛铁, 杨瑞东, 高军波, 毛家仁. 2015. 贵州织金寒武系磷矿沉积特征及灯影组古喀斯特面控矿特征研究. 地质学报, 89(12): 2374 ~ 2388.

全国地层委员会《中国地层表》编委会. 2014. 中国地层表(2014). 北京: 地质出版社.

魏国齐, 杨威, 杜金虎, 徐春春, 邹才能, 谢武仁, 曾富英, 武赛军. 2015. 四川盆地震旦纪一早寒武世克拉通内裂陷地质特征. 天然气工业, 35(1): 24 ~ 35.

汪泽成, 姜华, 王铜山, 鲁卫华, 谷志东, 徐安娜, 杨雨, 徐兆辉. 2014. 四川盆地桐湾期古地貌特征及成藏意义. 石油勘探与开发, 41(3): 305 ~ 312.

武赛军, 魏国齐, 杨威, 谢武仁, 曾富英. 2016. 四川盆地桐湾运动及其油气地质意义. 天然气地球科学, 27(01): 60 ~ 70.

王世谦, 陈更生, 黄先平. 2005. 四川盆地油气资源潜力及重点勘探领域. 石油学报, 26(增): 97 ~ 101.

邹才能, 杨智, 朱如凯, 张国生, 侯连华, 吴松涛, 陶士振, 袁选俊, 董大忠, 王玉满, 王岚, 黄金亮, 王淑芬. 2015. 中国非常规油气勘探开发与理论技术进展. 地质学报, 89(6): 979 ~ 1007.

朱日祥, 李献华, 侯先光, 潘永信, 王非, 邓成龙, 贺怀宇. 2009. 梅树村剖面离子探针锆石 U-Pb 年代学: 对前寒武纪—寒武纪界线的年代制约. 中国科学(D辑: 地球科学), 39(8): 1105 ~ 1111.

Brasier M, Cowie J and Taylor M. 1994. Decision on the Precambrian—Cambrian boundary stratotype. Episodes, 17(1): 3 ~ 8.

Condon D, Zhu Maoyan, Bowring S, Wang Wei, Yang Aihua and Jin Yuguan. 2005. U-Pb Ages from the Neoproterozoic Doushantuo Formation, China. Science, 308(1): 277 ~ 280.

Chen Daizhao, Zhou Xiqiang, Fu Yong, Wang Jianguo and Yan Detian. 2015. New U-Pb zircon ages of the Ediacaran—Cambrian boundary strata in South China. Terra Nova, 27(1): 62 ~ 68.

Chen Menge. 1982&. The new knowledge of the fossil assemblages from Maidiping section, Emei county, Sichuan with reference to the Simian—Cambrian boundary. Scientia Geologica Sinica, 17(3): 253 ~ 264.

Chen Nansheng, Yang Xiuzhen, Liu Dehan and Xiao Xuejun. 1982&. Lower Cambrian black argillaceous and arenaceous rock series in south China and its associated stratiform deposits. Mineral Deposits, 1(2): 39 ~ 51.

Compston W, Zhang Zichao, Cooper J A, Ma Guogan and Jenkins R J F. 2008. Further SHRIMP geochronology on the early Cambrian of

- South China. *American Journal of Science*, 308(4): 399~420.
- Chen Zhiming and Chen Qiyang. 1987#. Paleogeography of Yangzi platform and the characteristics of the phosphorite distribution of Early Meishucun stage, Early Cambrian. 22(3): 246~257.
- Deng Shenghui, Fan Ru, Li Xin, Zhang Shibin, Zhang Baomin and Lu Yuanzheng. 2015#. Subdivision and correlation of the Sinian (Ediacaran) system in the Sichuan basin and its adjacent area. *Journal of Stratigraphy*, 39(3): 239~254.
- Du Jinhu, Wang Zecheng, Zou Caineng, Xu Chunchun, Shen Ping, Zhang Baomin, Jiang Hua and Huang Shipeng. 2016#. Discovery of intra-cratonic rift in the Upper Yangtze and its control effect on the formation of Anyue giant gas field. *Acta Petrolei Sinica*, 37(1): 1~16.
- Du Jinhu, Zhou Caineng, Xu Chunchun, He Haiqing, Shen Ping, Yang Yueming, Li Yalin, Wei Guoqi, Wang Zecheng and Yang Yu. 2014#. Theoretical and technical innovations in strategic discovery of a giant gas field in Cambrian Longwangmiao Formation of central Sichuan paleo-uplift, Sichuan Basin. *Petroleum Exploration and Development*, 41(3): 268~277.
- Guo Qingjun, Shields G A, Liu Congqiang, Strauss H, Zhu Maoyan, Pi Daohui, Goldberg T and Yang Xinglian. 2007. Trace element chemostratigraphy of two Ediacaran—Cambrian successions in South China; Implications for organosedimentary metal enrichment and silicification in the Early Cambrian. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 254(1~2): 194~216.
- Griffin W L, Belousova E A, Shee S R, Pearson N J and Reilly S Y O. 2004. Archean crustal evolution in the northern Yilgarn Craton: U-Pb and Hf-isotope evidence from detrital zircons. *Precambrian Research*, 131(3~4): 231~282.
- Gu Zhidong, Di Xiufen and Yuan Miao. 2015#. Zircon U-Pb dating, geochemical characteristics and tectonic setting of basal granite in Sichuan Basin. *Acta Geologica Sinica*, 89(Supp): 296~296.
- He Hongchun and Qu Xiaoyan. 2015#. Present situation and prospect of shale gas exploration in Sichuan Basin. *Geological Review*, 61(Supp): 139~141.
- Jackson S E, Pearson N J, Griffin W L and Belousova E A. 2004. The application of laser ablation-inductively coupled plasma-mass spectrometry to in situ U-Pb zircon geochronology. *Chemical Geology*, 211(1~2): 47~69.
- Luo Beiwei, Wei Guoqi, Yang Wei and Dong Caiyuan. 2013a#. Study on the sea-level change during the Dengying stage in Sichuan Basin. *Geological Review*, 59(Supp): 221~222.
- Luo Beiwei, Wei Guoqi, Yang Wei and Dong Caiyuan. 2013b#. Reconstruction of the late Sinian paleo-ocean environment in Sichuan basin and its geological significance. *Geology in China*, 40(4): 1099~1111.
- Liu Jingjiang, Li Wei, Zhang Baomin, Zhou Hui, Yuan Xiaohong, Shan Xiuqin, Zhang Jing, Deng Shenghui, Gu Zhidong, Fan Ru, Wang Yongjun and Li Xin. 2015#. Sedimentary paleogeography of the Sinian in Upper Yangtze region. *Journal of Palaeogeography*, 17(6): 735~753.
- Ludwig K R. 2008. User's Manual for Isoplot 3.70: A Geochronological Toolkit for Microsoft Excel. Berkley Geochronology Center Special Publication, No. 4.
- Liu Pengju, Yin Chongyu, Chen Shouming, Li Meng, Gao Linzhi and Tang Feng. 2012#. Discussion on the Chronostratigraphic Subdivision of the Ediacaran (Sinian) in the Yangtze Gorges Area, South China. *Acta Geologica Sinica*, 86(6): 849~866.
- Li Zhongquan, Liu Ji, Li Ying, Hang Wenyan, Hong Haitao, Ying Danlin, Chen Xiao, Liu Ran, Duan Xinguo and Peng Jian. 2015#. Formation and evolution of Weiyan—Anyue extension—erosion groove in Sinian system, Sichuan Basin. *Petroleum Exploration and Development*, 42(1): 26~33.
- Li ZX, Li XH, Kinny P D, Wang J, Zhang S and Zhou H. 2003. Geochronology of Neoproterozoic syn-rift magmatism in the Yangtze Craton, South China and correlations with other continents: evidence for a mantle superplume that broke up Rodinia. *Precambrian Research*, 122(1~4): 85~109.
- Mou Nan and Wu Chaodong. 2005#. Characteristics and Phosphogenesis of Phosphorite of the Sinian—Cambrian, West Yangtze Area. *Acta Scientiarum Naturalium Universitatis Pekinensis*, 41(4): 551~562.
- Mao Tie, Yang Ruidong, Gao Junbo and Mao Jiaren. 2015#. Study of Sedimentary Feature of Cambrian Phosphorite and Ore - controlling Feature of Old Karst Surface of the Dengying Formation in Zhijin, Guizhou. *Acta Geologica Sinica*, 89(12): 2374~2388.
- National Commission on Stratigraphy of China. 2014#. The Stratigraphy Chart of China. Beijing: Geological Publishing House.
- Steiner M, Li Guoxiang, Qian Yi, Zhu Maoyan and Erdtmann B D. 2007. Neoproterozoic to Early Cambrian small shelly fossil assemblages and a revised biostratigraphic correlation of the Yangtze Platform (China). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 254(1~2): 67~99.
- Wei Guoqi, Yang Wei, Du Jinhu, Xu Chunchun, Zhou Caineng, Xie Wuren, Zeng Fuying and Wu Saijun. 2015#. Geological characteristics of the Sinian—Early Cambrian intracratonic rift, Sichuan Basin. *Natural Gas Industry*, 35(1): 24~35.
- Wu Saijun, Wei Guoqi, Yang Wei, Xie Wuren and Zeng Fuying. 2016#. Tongwan Movement and its geologic significances in Sichuan Basin. *Natural Gas Geoscience*, 27(1): 60~70.
- Wang Shiqian, Chen Gengsheng and Huang Xianping. 2005#. Petroleum resource potential and major exploration areas in Sichuan Basin. *Acta Petrolei Sinica*, 26(supp): 97~101.
- Wang Zecheng, Jiang Hua, Wang Tongshan, Lu Weihua, Gu Zhidong, Xu Anna, Yang Yu and Yang Zhaohui. 2014#. Paleogeomorphology formed during Tongwan tectonization in Sichuan Basin and its significance for hydrocarbon accumulation. *Petroleum Exploration and Development*, 41(3): 305~312.
- Xu Lingang, Lehmann B, Mao Jingwen, Qu Wenjun and Du Andao. 2011. Re-Os Age of polymetallic Ni—Mo—PGE—Au mineralization in early Cambrian black shales of south China—A reassessment. *Economic Geology*, 106: 511~522.
- Zou Caineng, Yang Zhi, Zhu Rukai, Zhang Guosheng, Hou Lianhua, Wu Songtao, Tao Shizhen, Yuan Xuanjun, Dong Dazhong, Wang Yuman, Wang lan, Huang Jinliang and Wang Shufen. 2015#. Progress in China's Unconventional Oil & Gas Exploration and Development and Theoretical Technologies. *Acta Geologica Sinica*, 89(6): 979~1007.
- Zhuo Jiewen, Jiang Zhuofei, Jiang Xinsheng, Wang Jian, Cai Juanjuan, Xiong Guoqing, Lu Junze, Cui Xiaozhuang and Liu Jianhui. 2017#. SHRIMP zircon U-Pb ages for the stratotype section of Neoproterozoic Suxiong Formation in western Sichuan province and their geological significance. *Geological Review*, 63(1): 177~188.
- Zhu Maoyan, Zhang Junming, Steiner M, Yang Aihua, Li Guoxiang and Erdtmann B D. 2003. Sinian—Cambrian stratigraphic framework for shallow- to deep-water environments of the Yangtze Platform; An

integrated approach. *Progress in Natural Science*, 13(12): 951 ~ 960.

Zhu Maoyan. 2007. From snowball earth to the Cambrian bioradiation: calibration of Ediacaran—Cambrian earth history in South China. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 254: 1 ~ 6.

Zhu Rixiang, Li Xianhua, Hou Xiangang, Pan Yongxin, Wang Fei, Deng Chenglong and He Huaiyu. 2009. SIMS U-Pb zircon age of a tuff layer in the Meishucun section, Yunnan, southwest China: Constraint on the age of the Precambrian—Cambrian boundary. *Science in China (Series D)*, 39(8): 1105 ~ 1111.

LA-ICP-MS U-Pb Zircon Ages of Volcaniclastic Beds of the Third Member of the Sinian (Ediacaran) Dengying Formation in Leshan, Sichuan, and a Discussion on the Rift Evolution in the Basin

ZI Jinping¹⁾, JIA Dong¹⁾, WEI Guoqi²⁾, YANG Zhenyu³⁾, ZHANG Yong¹⁾,
HU Jing¹⁾, SHEN Shuxin¹⁾

1) State Key Laboratory for Mineral Deposit Research, Nanjing University, Nanjing, 210023;

2) Research Institute of Petroleum Exploration and Development, Langfang Branch, PetroChina, Langfang, Hebei, 065007;

3) College of Resources, Environment and Tourism, Capital Normal University, Beijing, 100048

Abstract: The late-Sinian (Ediacaran) to early-Cambrian intracratonic rift in central Sichuan basin has a close relationship with the formation of Anyue Sinian (Ediacaran)—Cambrian giant gas field. geochronology study of the Dengying Formation is significant in understanding the evolution of the intracratonic rift. LA-ICP-MS U-Pb analysis of volcanic ash beds in the Third Member of the Dengying Formation at the Xianfeng Section, Sichuan, Southwest China yield a highly reliable Concordia age of 539.6 ± 1.4 Ma (MSWD = 0.22), suggesting a lower placement of the Precambrian—Cambrian boundary in the Sichuan Basin. We established a evolution time frame of this intracratonic rift based on four isochronous strata, the bottom of the Dengying Formation (551.1 Ma), the Third Member of Dengying Formation (539.6 Ma), the Maidiping Formation (535.2 Ma) and the Ni—Mo sulfide enriched layer in the lower Cambrian (521 Ma). The boundary of the Maidiping Formation and the Dengying Formation inside and outside the rift is not synchronous. According to those, we propose that the formation of the late-Sinian (Ediacaran) to early-Cambrian intracratonic rift in central Sichuan Basin was mainly controlled by the boundary fault. Due to the deep-water depositional environment in the rift, the Maidiping Formation began to deposit rapidly after the deposition of the thin Third and Fourth Member of the Dengying Formation, the Maidiping Formation and the Qiongzusi Formation inside the rift and the Fourth Member of the Dengying Formation outside the rift are contemporary heteropical sediments.

Keywords: Sichuan Basin; Cambrian; Sinian (Ediacaran); the Dengying Formation; U-Pb geochronology; intracratonic rift

Acknowledgements: This study was financially supported by the National Science and Technology Major Projects (No. 2016ZX05003-001 and No. 2017ZX05008-001) and the Natural Science Foundation of China (No. 41230208).

First author: ZI Jinping, Male, Born in Hengyang city, Hunan Province. Now studying at School of Earth Sciences and Engineering, Nanjing University for a postgraduate degree, Major is structural geology. Email: zijinping@qq.com

Corresponding author: JIA Dong, Professor of School of Earth Sciences and Engineering, Nanjing University, Email: djia@nju.edu.cn

Manuscript received on: 2016-12-19; Accepted on: 2017-05-10; Edited by: LIU Zhiqiang

Doi: 10.16509/j.georeview.2017.04.014