

四川盆地下古生界硅质页岩层系中硅质来源及其对有机质保存的影响初探



www.
geojournals.cn/georev

谢小敏^{1,2,3)}, 刘伟新³⁾, 张瀛⁴⁾, 赵迪斐⁵⁾, 唐友军¹⁾, 申宝剑³⁾, 李志明³⁾

1) 油气资源与勘探技术教育部重点实验室, 长江大学资源与环境学院, 武汉, 430100;

2) 南京宏创地质勘查技术服务有限公司, 南京, 211111;

3) 中石化石油勘探开发研究院无锡石油地质研究所, 江苏无锡, 214126;

4) 重庆科技学院, 重庆, 401331;

5) 中国矿业大学煤层气资源与成藏过程教育部重点实验室, 江苏徐州, 221008

内容提要:四川盆地下古生界两套硅质页岩层系(寒武系底部页岩和奥陶系—志留系五峰组—龙马溪组页岩)是其重要的页岩气产层,五峰组—龙马溪组硅质含量与 *TOC* 及含气量呈正相关性,暗示硅质矿物对该地区页岩气的生成具有重要意义。但硅质来源复杂,且存在后期硅质流体的注入。因此,本文在总结前人的研究成果基础上,对四川盆地寒武系底部硅质岩系和五峰组—龙马溪组硅质岩系的硅质特征进行了详细的岩石学分析,揭示了3种硅质来源特征:①同沉积无机硅质流体;②生物硅;③后期无机硅质流体。其中寒武系底部以同沉积无机硅质流体和生物硅来源为主,同沉积硅质流体导致有机质快速石化埋藏,有机质内部结构及形态保存较好;五峰组—龙马溪组硅质来源以后期无机硅质流体和生物硅来源为主,有机质在保存过程中受硅质流体的影响很小,有机质腐泥化作用充分,内部结构和形态多数保存较差。该研究初步揭示了两套页岩层系的硅质发育岩石学特征,及其对有机质保存的影响,期望能为四川盆地两套页岩气储层差异性提供新的思路 and 基础地质数据。

关键词:硅质页岩;硅质来源;有机质保存;下古生界;四川盆地

页岩气资源,是自生自储于富有机质页岩层系中的非常规天然气资源(Jarvie et al., 2007; Loucks and Ruppel, 2007)。四川盆地及其周缘是中国最重要的页岩气产区,具有页岩气地质储量为 $62.56 \times 10^{12} \text{ m}^3$,可采储量 $9.94 \times 10^{12} \text{ m}^3$ (Zhang Yuying et al., 2019)。该地区下古生界发育两套重要的页岩气储层,下寒武统牛蹄塘组或筇竹寺组(孙玮等, 2012; 谢小敏等, 2015; 周泽等, 2016; Leng Jigao et al., 2016; Liu Honglin et al., 2016; Wang Ruyue et al., 2016; Zeng Weite et al., 2016; Zhang Yuying et al., 2018; 冯明友等, 2018; 吴安彬等, 2020)和上奥陶统五峰组—下志留统龙马溪组(金之钧等, 2016; 冯动军等, 2016; Li Ang et al., 2017; Shao Xinhe et al., 2017; 魏祥峰等, 2017)。前期研究表明,龙马溪组页岩气优质储层通常位于该套地层底部富含硅质页岩层段,其自然伽马值较高($>150 \text{ API}$),岩石有机碳含量高($>2\%$),含气量高($>2 \text{ m}^3/\text{t}$; 王玉满

等, 2012; 刘树根等, 2011, 2013; 郭彤楼和张汉荣, 2014; 赵佩等, 2014; 毕赫等, 2014)。以典型的焦页1页岩气井为例,五峰组—龙马溪组下段页岩 *TOC* 含量为 $0.55\% \sim 5.89\%$, 平均 2.5% , 其中底部硅质页岩 *TOC* 含量最高($2.56\% \sim 4.97\%$, 平均 3.89%), 是该井页岩气的主要产气层段(金之钧等, 2016)。前人的研究发现,涪陵地区五峰组—龙马溪组下部优质页岩硅质矿物含量与 *TOC* 含量具有较好的正相关性,硅质矿物含量越高,有机碳含量越高(金之钧等, 2016; Yang Feng et al., 2016; Xu Zhuang et al., 2019);同时岩石孔隙度与硅质矿物之间也具有很好的正相关性(Yang Feng et al., 2016),暗示硅质组成对该地区页岩气生成具有潜在的重要意义。

吴蓝宇等(2016)对涪陵地区典型五峰组—龙马溪组页岩气井研究发现,这套硅质页岩发育地层有机碳含量与含气量之间具有很好的相关性($R^2 =$

注:本文为国家自然科学基金委员会面上自然基金资助项目(编号:41972163)的成果。

收稿日期:2020-06-14; 改回日期:2021-01-11; 网络首发:2021-02-20; 责任编辑:刘志强。Doi: 10.16509/j.georeview.2021.02.012

作者简介:谢小敏,女,1984年生,博士,现在长江大学主要从事岩石学和地球化学综合研究工作; Email: xiaominxie2019@sina.com。




图1 四川盆地涪陵气田五峰组—龙马溪组硅质页岩层段页岩含气量与 TOC (a, 据吴蓝宇等, 2016 修改) 和 TOC 与 SiO_2 含量 (b, 据 Xu Zhuang et al., 2019 修改) 相关散点图

Fig. 1 Relationships of gas content vs TOC (a, modified from Wu Lanyu et al., 2016) and TOC vs SiO_2 (b, modified from Xu Zhuang et al., 2019) in the shale members of the Wufeng Formation—Longmaxi Formation in Fuling gas field, Sichuan Basin

0.6671, 图 1a)。同时, 该层段 TOC 含量与 SiO_2 含量之间具有很好的相关性 (图 1b, Xu Zhuang et al., 2019)。而非优质页岩段硅质矿物含量与 TOC 含量相关性较差, 揭示了优质页岩段与非优质页岩段硅质矿物成因的不同 (金之钧等, 2016)。因此, 五峰组—龙马溪组硅质页岩储层中硅质含量与 TOC 及含气量呈正相关性, 暗示硅质矿物对该地区页岩气的生成具有重要意义。

但由于硅质来源和成因较复杂, 有学者对该地区下古生界硅质岩的成因也做了详细的研究, 并计算了过剩硅 (excess silicon) 的含量 (卢龙飞等, 2016; 刘江涛等, 2017; Zhang Kun et al., 2018)。但实际上, 过剩硅是指高于正常碎屑沉积环境下的 SiO_2 含量, 并不能区分硅质来源的生物硅和热液硅特征, 而热液硅也存在同沉积与后期注入的不同。因此, 本文在前人研究的基础上, 采集了贵州羊跳地区寒武系底部牛蹄塘组野外剖面样品、四川盆地焦石坝和丁山地区五峰组—龙马溪组岩芯样品, 进行了显微镜下岩石学特征的精细观察。提出了三种可能的硅质来源, 并初步探讨了两套硅质页岩层系硅质来源的不同及其对有机质保存的影响, 期望能为四川盆地两套页岩气储层差异性提供新的思路和基础地质数据。

1 关于硅质岩的成因

硅质的成岩作用主要有化学作用、生物和生物化学作用及火山作用、热水作用 (张亚冠等, 2015)。前人对沉积岩中的硅质进行了岩石学、常量元素、微

量元素、稀土元素、同位素分布特征的研究, 揭示了硅质的成因认识。目前有 3 种主要的成因认识:

(1) 热液成因: 在常温下海水中的硅含量极低 ($<10 \times 10^{-6}$; Calvert, 1983), 不可能达到饱和而直接沉淀, 仅靠放射虫和海绵等硅质生物形成厚度很大的硅质沉积是很难的 (涂光炽, 1990)。而硅的溶解度随着水温的增高而迅速增加, 海水中硅的含量在 150°C 时达到 600×10^{-6} (Krauskopf, 1956), 在 200°C 时海水中的含硅量是 50°C 的 10 倍 (Holland, 1967)。富硅的热液遇到冷海水后, 硅的含量可能超过常温水中溶解度的 10~20 倍 (Rona, 1978), 呈过饱和状态。因此, 在海底热泉附近可能有硅质的直接沉淀 (Rona et al., 1980)。热泉附近因营养丰富, 硅质生物及其他生物十分繁荣, Dick (2019) 在现代东太平洋海底热液喷口黑烟柱附近观察到了大量的微生物和底栖动物。20 世纪 80 年代以来, 国内外很多专家学者对硅质岩研究表明, 海底或大陆许多硅质岩为热液成因硅质岩 (Adachi et al., 1986; Yamamoto, 1987; 陈洪德和曾允孚, 1989; 伊海生等, 1989; 周永章, 1990; 周永章等, 1994; 夏邦栋等, 1995; 冯胜斌等, 2007; 李凤杰等, 2010)。寒武系底部硅质岩发育时期, 存在大规模的火山活动和海底喷流作用; 这为硅质岩的热液来源提供了必要的条件, 同时也导致了寒武纪早期海洋的物质循环和水体化学性质的特殊性 (Steiner et al., 2001; Kamber and Weeb, 2001; Chen Daizhao et al., 2009)。Adachi 等 (1986) 和 Yamamoto (1987) 认为热液中形成的硅质具有 SiO_2 含量高, Al_2O_3 和 TiO_2 含量低的

特征。Bostrom (1983) 认为与热液沉积有关沉积物中,富 Fe、Mn, 贫 Cu、Co 和 Ni; 而 Al 的富集则与陆源物质输入有关。因此, $Al/(Al+Fe+Mn)$ 值可作为衡量沉积物中热液沉积物含量多少的标志 (Bostrom et al., 1969; Adachi et al., 1986; Yamamoto, 1987); $(Cu+Co+Ni)-Fe-Mn$ 三角图也用于区分热液与非热液沉积 (Rona, 1988)。

(2) 生物成因: 硅质主要来源于硅质生物, 如海绵骨针、放射虫、硅鞭藻、硅藻等 (Boggs, 2006; Robert and Stephen, 2007; Day-Stirrat et al., 2010; Milliken et al., 2012)。五峰组—龙马溪组产气层段硅质页岩中石英含量高, 一般大于 45%, 认为这类硅质主要来源于生物硅 (杨玉卿和冯增昭, 1997; Schieber et al., 2000; Loucks and Ruppel, 2007; 杨水源和姚静, 2008; 卢龙飞等, 2016)。Yang Xiangrong 等 (2018) 分析了硅质岩中的石英含量与生物钡含量呈正相关, 认为生物硅质是该地区重要的硅质的来源。王玉满等 (2016) 对川南龙马溪组页岩岩相进行了划分, 认为硅质页岩 (华蓥—长宁一带) 和钙质硅质混合页岩 (威远—泸州—綦江以南 (王玉满等, 2015) 是深水沉积环境的特有岩相, 发育水深在 100 m 以下, 富含有机质, 古生物化石, 如放射虫、海绵骨针、尖笔石等 (刘伟等, 2010; 樊隽轩等, 2012, 2013; 张春明等, 2012; 王淑芳等, 2014)。在早寒武世硅质岩中也发现了大量的放射虫和海绵骨针等硅质生物 (张位华等, 2003; 谢小敏等, 2015)。因此, 硅质生物是这两套页岩储层硅质的一个重要来源。

(3) 交代成因: 交代成因硅质是指在岩石沉积之后, 硅质流体交代原生矿物形成的硅质矿物。一般来说, 交代成因硅质岩的原岩多为碳酸盐岩, 有成岩期交代 (杨吉和李英, 2002) 和地表次生交代等多种类型 (李文恒, 1980)。朱嗣昭 (1999) 在南京及巢湖地区下二叠统孤峰组中发现了这类硅质岩 (称为“浮石状隧石”), 是一种交代碳酸盐岩成因的硅质岩, 显著特征是岩石中存在大量菱形和不规则状孔隙, 孔隙周边和孔隙间主要由微晶石英组成。施贵军等 (1999) 在皖南浙西交界地区黄龙组也发现交代成因硅质岩。郭福生等 (2003) 对浙江江山晚石炭世藕塘组上段发现“钙骨假象燧石岩”, 特征是大量钙质生物碎屑 (包括腕足类、海百合茎、珊瑚等) 被完全硅化, 常见由多个微晶石英交代的自形细晶白云石假象以及碳酸盐交代残余; 元素 $Al/(Al+Fe+Mn)$ 值显示, 该地区硅质岩全部属于正常海水

沉积区, 微量元素和稀土元素特征等显示该硅质源于与热水沉积无密切关系, 属于成岩交代成因硅质岩。Marin-Carbonne 等 (2011) 对南非 Onverwacht Group 的燧石进行综合的原位氧同位素和硅同位素, 及硅质脉体中流体包裹体特征, 认为该燧石中硅质为交代成因。Rakociński 和 Boreuch (2016) 在波兰晚泥盆弗拉斯阶碳酸盐岩中也发现了硅化的头足类动物化石, 认为快速的硅化有利于化石带的保存。

实际地质样品中, 硅质的成因往往是综合的, 如生物和热水综合成因 (田云涛等, 2007) 等。据 Clayton (1986) 和 Douthitt (1982) 等的研究, 不同来源硅质具有不同的 $\delta^{30}Si$ 值, 从低温地下水中自生沉淀的石英 $\delta^{30}Si$ 值 ($>1.4‰$) $>$ 成岩过程中次生加大石英的 $\delta^{30}Si$ 值 ($0.8‰ \sim 1.4‰$) $>$ 热水来源石英的 $\delta^{30}Si$ 值 ($-1.5‰ \sim 0.8‰$)。宋天锐和丁梯平 (1989) 分析认为, 生物成因硅质岩的 $\delta^{30}Si$ 值为 $-1.1‰ \sim -1.7‰$, 交代成因硅质岩的 $\delta^{30}Si$ 值为 $2.4‰ \sim 3.4‰$, 火山喷发—化学沉积硅质岩的 $\delta^{30}Si$ 值为 $-0.4‰ \sim -0.5‰$ 。此外, 常量元素组成、微量元素组成、稀土、放射性同位素 (Rb-Sr、K-Ar、Pb 等)、稳定同位素 (H、C、O、 $^{87}Sr/^{86}Sr$ 、Sm-Nd、Si 等)、稀有气体元素 (Ar) 和有机地球化学等方面对不同成因硅质进行了精细的研究工作 (宋天锐和丁梯平, 1989; Murray, 1990, 1992; Murray et al., 1991; 盛吉虎和陈中新, 1998), 获得了不同成因硅质来源的地球化学信息。Holdaway 和 Clayton (1982) 年提出过量硅 (excess silicon) 的概念, 是指高于正常碎屑沉积环境下的 SiO_2 含量, 其计算公式为

$$Si_{\text{excess}} = Si_{\text{sample}} - (Si/Al)_{\text{background}} \times Al_{\text{sample}},$$

其中 $(Si/Al)_{\text{background}}$ 采用平均页岩比值 3.11, 该方法广泛应用于硅质矿物研究中 (刘江涛等, 2017; Zhang Kun et al., 2018), 但对于不同成因非碎屑来源的硅质的相对含量定量计算分析还比较缺乏。

2 寒武系底部硅质岩石学特征及其对有机质保存的影响

寒武系底部与五峰组—龙马溪组硅质页岩储层相比, 寒武系底部硅质页岩具有有机碳含量和有机质热演化程度更高 (Tan Jingqiang et al., 2014; Wang Chao et al., 2018; Zhao Jianhua et al., 2018; Zhang Yuying et al., 2018, 2019), 页岩储层厚度更大, 脆性矿物 (以硅质为主) 更高 (平均含量接近 80%; Zhang Yuying et al., 2019) 等特点。然而, 该层位页岩气的勘探开发并未取得很大的突破。虽然有学者




图2 四川盆地边缘早寒武世牛蹄塘组底部硅质页岩显微照片:(a)—(c) 硅质页岩中保存较好的生物化石,(a)为海绵骨针,(b)为疑源类,(c)为须腕动物;(d)—(f) 具十字消光玉髓,(e)和(f)是(d)中玉髓的局部放大;(a)—(e)为透射光下偏光显微照片;(f) 为透射光正交偏光显微照片

Fig. 2 Micrographs of siliceous shales from Lower Cambrian Niutitang Formation in Sichuan Basin: (a)—(c) Well-preserved fossils in siliceous shales, (a) sponge spicule, (b) acritarch, (c) pogonophora; (d)—(f) chalcedony with cross extinction; (a)—(e) are micrographs of polarized transmitted light; (f) is micrograph of orthogonal transmitted light

将其归咎于寒武系页岩经历了更复杂的构造活动改造,裂缝发育,导致后期保存条件较差(Zhang Yuying et al., 2019)。但实际上,两套硅质页岩储层的硅质组成不同,也可能是其原因之一。

在四川盆地边缘贵州寒武系底部牛蹄塘组硅质页岩层系中,硅质岩呈层状、透镜状产出(谢小敏等,2015),显微镜下硅质透光下呈圆形,正交光下具十字消光(图2),与美国黄石公园硅质热泉沉淀特征类似(Guidry et al., 2003)。谢小敏等(2015)在贵州寒武系地层中也见到了玉髓化现象和典型的热液生物,即须腕动物蠕虫(Monika et al., 2005; Kenneth, 2005; 梁狄刚等,2009),从而认为该套硅质岩形成与海底热液活动有关。寒武纪早期是大陆内部强烈拉张作用引起强烈的热液作用时期,前人通过微量元素分析、稀土元素、同位素地球化学、岩石学和矿物学分析,已做了大量精细的研究确定了这套地层沉积时期受到海底热液的影响(李胜荣等,1995; Steiner et al., 2001; Jiang Shaoyong et al., 2003; 韩善楚等,2003; 杨剑等,2004; 杨兴莲等,2007; 杨瑞东等,2007; 谢小敏等,2015; Liu Zhanhong et al., 2015)。

在这样一套硅质页岩层系中,发现了很多生物形态保存较好的原生有机质,如海绵骨针、疑源类、须腕动物及其他藻类等(杨瑞东等,2005; 杨兴莲等,2010; 谢小敏等,2015)。有机质保存完好的部分都是在富含硅质的区域(图2a~c),如疑源类表面微小的纹饰都能清晰可见(图2b);且疑源类细胞呈圆形,未出现沉积有机质受细菌降解、成岩压实引起的细胞变形。一般来说,油气生成过程中,有机质在复杂的沉积—埋藏过程中会经历细菌腐泥化作用,后续成岩—生烃演化阶段,生烃作用也会改造有机质形态结构,最终使得有机质的形态保存较差,尤其是生成大量油气之后,形态保存较好的有机质就更少。因此,在传统显微镜和在高精度的扫描电镜下,能鉴定出来的形态特征保存较好的生物,是否就代表了烃源岩中主力的生烃有机质特征,存在较大的疑议(Hutton, 1994)。初步推测这些形态保存完好的生物有机质可能与同沉积的热液事件有关,热液的加入使得生物快速死亡—埋藏—保存在硅质岩层中。也就是说,同沉积热液硅质可能瞬间石化了有机质,从而很好的保护了有机质的外形,这类现象在马达加斯加的浅海海岸海洋玛瑙中比较常见(图




图 3 马达加斯加的浅海海岸海洋玛瑙显微照片,显示玉髓沉淀过程中包裹了有机质(OM)

Fig. 3 Micrographs of marine agate from the shallow coast of Madagascar, showing the inclusion of organic matter (OM) during the deposition of chalcedony

3)。这些保存较好的有机质一般发现在硅质含量较高的含泥或泥质硅质岩中,在这类岩石中,能指示成烃过程的固体沥青的含量并不高。而在硅质页岩或页岩中,有机质的保存相对较差,明显经历了沉积成岩及细菌的改造作用。这类快速被保存的有机质,由于并未经历缓慢的细菌腐泥化作用,后期埋藏过程中又受到硬度较大的硅质矿物的保护。

3 五峰组—龙马溪组硅质岩石学特征及其对有机质保存的影响

四川盆地五峰组—龙马溪组底部硅质页岩中硅质特征则完全不同,虽然有学者在该套硅质页岩中发现了海绵骨针和放射虫(卢龙飞等,2016;刘江涛等,2017),认为其主要是生物来源的。实际上,与寒武系底部的硅质页岩相比,五峰组—龙马溪组底部硅质页岩硅质生物的发育程度要少很多,且过量硅含量为 0~20%,而寒武系的为 20%~50% (Zhang Kun et al., 2018)。Yang Xiangrong 等(2018) 对华南龙马溪组底部硅质页岩样品分析,发现 TOC 与硅质呈正相关关系时,可能与生物硅来源有关;但 TOC 大于 3.2 %时,TOC 与硅质含量呈现负相关关系($R^2=0.88$,图 4)。是否可能在 TOC 高于 3.2%的岩层中,硅质来源存在差异?




图 4 四川盆地志留系龙马溪组硅质页岩和黏土质页岩有机碳含量 (TOC) 与 SiO₂ 含量相关关系图 (据 Yang Rongxiang et al., 2018 修改)

Fig. 4 The plots of TOC vs SiO₂ contents of siliceous and argillaceous shale in Sichuan Basin (modified from Yang Rongxiang et al., 2018)

对丁山地区该套硅质页岩底部的薄片显微特征分析(图 5a—d),发现较多硅质脉体,有的硅质脉体切断了方解石脉,或出现方解石脉被硅质脉体交代残余(图 5a、b),硅质颗粒边缘还残留少量钙质残余(图 5c),笔石皮层被硅化(图 5d)等现象,表明硅




图5 四川盆地丁山地区(a)—(d)和焦石坝地区(e)—(f)五峰组—龙马溪组底部硅质页岩中硅质正交光下显微照片:(a)、(b)方解石脉被硅质条带交代残余;(c)硅质颗粒边缘发现钙质物质残留;(d)被硅化的笔石皮层;(e)、(f)焦石坝地区五峰组—龙马溪组底部硅质页岩中,发育玉髓包裹有机质(OM)现象

Fig. 5 Microphotos under orthogonal light of siliceous shales of the lower Wufeng Formation—the Longmaxi Formation in Dingshan area (a) —(d) and Jiaoshiba area (e)、(f): (a)、(b) calcite veins are metasomatic with silica bands; (c) calcium residues were found at the edge of the silica particles; (e)、(f) the organic matter (OM) were found to be included by silica in Jiaoshiba area

质存在交代成因,切割关系说明硅质脉体充注的时间晚于方解石脉形成时间。在焦石坝地区,五峰组—龙马溪组底部见到较多的硅质包裹有机质的特征(图5e、f),这类硅质正交下呈十字消光(玉髓现象),有机质的形态大小不一,棱角清晰,有机岩石学特征显示可能为固体沥青。也就是说,硅质可能是在油气生成的高峰时期注入,包裹了部分油(后期演化为固体沥青),这样形成了形态各异的硅质+有机质组合。这套硅质页岩中,除了笔石有机质保存较好外,藻类有机质主要呈无定形体保存,显示有机质在沉积埋藏过程中经历了充分的腐泥化过程,有助于后期油气的生成,晚期注入的硅质在对原生有机质的生烃潜力上影响不大。初步的电子探针分析,焦石坝地区硅质较丁山地区硅质脉体富含Fe元素,显示不同的硅质流体充注特征。

4 结论

对寒武系底部硅质岩和五峰龙马溪硅质岩的特征分析揭示了3种硅质来源特征:①同沉积无机硅质流体;②生物硅;③后期无机硅质流体。虽然两套

储层均存在硅质成分,但是硅质特征,来源及形成时间不同。其中寒武系底部以同沉积无机硅质流体和生物硅来源为主,同沉积硅质流体导致有机质快速石化埋藏,有机质(如凝源类等)内部结构及形态保存较好;五峰龙马溪组硅质来源以后期无机硅质流体和生物硅来源为主,有机质在保存过程中受硅质流体的影响很小,有机质腐泥化作用充分,内部结构和形态多数保存较差,以无定形体为主。因此,后期研究有必要将硅质岩的成分及来源进行精细分析,充分考虑硅质的岩石学特征及形成环境,区分同沉积时期的硅质与后期硅质矿物特征及含量,结合有机质含量、页岩储层特征进行相关性分析,探讨硅质成因对原生有机质和次生固体沥青有机质的影响,进而探讨其对页岩气生成的影响。

致谢:感谢边立曾、曹剑和高建峰教授对文章初稿提出了宝贵的意见,感谢林静文和吴芬婷在文章修改过程中提供的帮助,两位评审专家和责任编辑对本文提出了很多宝贵的修改意见和建议,在此深表谢意!

参 考 文 献 / References

(The literature whose publishing year followed by a “&” is in Chinese with English abstract; The literature whose publishing year followed by a “#” is in Chinese without English abstract)

毕赫, 姜振学, 李鹏, 李卓, 唐相路, 张定宇, 许野. 2014. 渝东南地区黔江凹陷五峰组—龙马溪组页岩储层特征及其对含气量的影响. 天然气地球科学, 25(8): 1275~1283.

陈洪德, 曾允孚. 1989. 广西丹池盆地上泥盆统榴江组硅质岩沉积特征及成因讨论. 矿物岩石, 9(4): 22~28.

韩善楚, 胡凯, 曹剑. 2013. 华南早寒武世黑色岩系重晶石矿床环带钼冰长石新发现及其意义. 地质论评, 59(6): 1143~1149.

樊隽轩, 吴磊, 陈中阳, 陈清, 张琳娜, Soo K Y, Singh N, 李学康, 王媛. 2013. 四川兴文县麒麟乡五峰组—龙马溪组黑色页岩的生物地层序列. 地层学杂志, 37(4): 513~520.

樊隽轩, Melchin M J, 陈旭, 王恽, 张元动, 陈清, 迟昭利, 陈峰. 2012. 华南奥陶—志留系龙马溪组黑色笔石页岩的生物地层学. 中国科学: 地球科学, 42(1): 130~139.

冯动军, 胡宗全, 高波, 彭勇民, 杜伟. 2016. 川东南地区五峰组—龙马溪组页岩气成藏条件分析. 地质论评, 62(6): 1521~1532.

冯胜斌, 周洪瑞, 燕长海, 彭翼, 袁效奇, 贺静. 2007. 东秦岭二郎坪群硅质岩热水沉积地球化学特征及其地质意义. 沉积学报, 25(2): 564~573.

冯明友, 郑江, 刘田, 刘小洪, 张家强, 梅文华, 孙霓源, 龙涛. 2018. 川北地区寒武系筇竹寺组钙质结核裂缝充填物特征及指示意义. 地质论评, 64(3): 711~722.

郭福生, 林子瑜, 杜杨松, 刘林清, 严兆彬. 2003. 一种特殊类型硅质岩的特征与成因研究. 地学前缘, 10(4): 573~581.

郭彤楼, 张汉荣. 2014. 四川盆地焦石坝页岩气田形成于富集高产模式. 石油勘探与开发, 41(1): 28~36.

金之钧, 胡宗全, 高波, 赵建华. 2016. 川东南地区五峰组—龙马溪组页岩气富集与高产控制因素. 地学前缘, 23(1): 1~9.

李凤杰, 刘殿鹤, 刘琪. 2010. 四川宣汉地区吴家坪组硅质岩地球化学特征及其成因探讨. 天然气地球科学, 21(1): 62~67.

李胜荣, 高振敏. 1995. 湘黔地区牛蹄塘组黑色岩系稀土特征:——兼论海相热水沉积岩稀土模式. 海相油气地质, 14(2): 1~19.

李文恒. 1980. 一种硅质岩成因的初步探讨. 地质论评, 26(5): 442~445.

梁狄刚, 郭彤楼, 边立曾, 陈建平, 赵醇. 2009. 南方四套区域性海相烃源岩的沉积相及发育的控制因素. 海相油气地质, 14(2): 1~19.

刘江涛, 李永杰, 张元春, 刘双莲, 蔡英杰. 2017. 焦石坝五峰组—龙马溪组页岩硅质生物成因的证据及其地质意义. 中国石油大学学报(自然科学版), 41(1): 34~41.

刘树根, 马文辛, Luba J, 黄文, 曾祥亮, 张长俊. 2011. 四川盆地东部地区下志留统龙马溪组页岩储层特征. 岩石学报, 27(8): 2239~2252.

刘树根, 王世玉, 孙玮, 冉波, 杨迪, 罗超, 叶玥豪, 白志强, 邱嘉文. 2013. 四川盆地及其周缘五峰组—龙马溪组黑色页岩特征. 成都理工大学学报: 自然科学版, 40(6): 621~639.

刘伟, 许效松, 冯心涛, 孙媛媛. 2010. 中上扬子上奥陶统五峰组含放射虫硅质岩与古环境. 沉积与特提斯地质, 30(3): 65~70.

卢龙飞, 秦建中, 申宝剑, 腾格尔, 刘伟新, 张庆珍. 2016. 川东南涪陵地区五峰—龙马溪组硅质页岩的生物成因成因及其油气地质意义. 石油实验地质, 38(4): 460~466.

盛吉虎, 陈中新. 1998. 硅质岩沉积地球化学研究现状. 地域研究

与开发, 17(增刊): 117~121.

施贵军, 杨湘宁. 1999. 皖南浙西交界地区黄龙组的硅质灰岩. 高校地质学报, 5(1): 113~114.

宋天锐, 丁梯平. 1989. 硅质岩中的硅同位素($\delta^{30}\text{Si}$)应用于沉积相分析的新尝试. 科学通报, 34(18): 1408~1411.

孙玮, 刘树根, 冉波, 王世玉, 叶玥豪, 罗超, 田梦娜. 2012. 四川盆地及周缘地区牛蹄塘组页岩气概况及前景评价. 成都理工大学学报(自然科学版), 39(2): 170~175.

田云涛, 冯庆来, 李琴. 2007. 桂西南柳桥地区上二叠统大隆组层状硅质岩成因和沉积环境. 沉积学报, 25(5): 671~677.

涂光炽. 1990. 本世纪 80 年代地球科学若干问题进展. 地质评论, 36(6): 510~517.

王淑芳, 邹才能, 董大忠, 王玉满, 黄金亮, 郭召杰. 2014. 四川盆地富有机质页岩硅质生物成因及对页岩气开发的意义. 北京大学学报: 自然科学版, 50(3): 476~486.

王玉满, 董大忠, 李建忠, 王社教, 李新景, 王黎, 程克明, 黄金亮. 2012. 川南下志留统龙马溪组页岩气储层特征. 石油学报, 33(4): 551~561.

王玉满, 董大忠, 李新景, 黄金亮, 王淑芳, 吴伟. 2015. 四川盆地及周边下志留统龙马溪组地层层序与沉积特征. 天然气工业, 35(3): 12~21.

王玉满, 王淑芳, 董大忠, 李新晨, 黄金亮, 张晨晨, 管全中. 2016. 川南下志留统龙马溪组页岩岩相表征. 地学前缘, 23(1): 119~133.

魏祥峰, 赵正宝, 王庆波, 刘珠江, 周敏, 张晖. 2017. 川东南綦江丁山地区上奥陶统五峰组一下志留统龙马溪组页岩气地质条件综合评价. 地质论评, 63(1): 153~164.

吴蓝宇, 胡东风, 陆永潮, 刘若冰, 刘晓峰. 2016. 四川盆地涪陵气田五峰组—龙马溪组页岩优势岩相. 石油勘探与开发, 43(2): 189~197.

吴安彬, 张景坤, 王井伶, 罗家国, 罗群, 姜振学. 2020. 富有机质页岩方解石脉成因、成岩模式与地质意义——以四川盆地南部龙马溪组为例. 地质论评, 66(1): 88~99.

夏邦栋, 钟立荣, 方中, 吕洪波. 1995. 下扬子区早二叠世孤峰组层状硅质岩成因. 地质学报, 69(2): 125~137.

谢小敏, 腾格尔, 秦建中, 张庆珍, 边立曾, 尹磊明. 2015. 贵州凯里寒武系底部硅质岩系生物组成、沉积环境与烃源岩发育关系研究. 地质学报, 89(2): 425~439.

杨剑, 易发称, 侯兰杰. 2004. 黔北黑色岩系的岩石地球化学特征和成因. 矿物学报, 24(3): 285~289.

杨吉, 李英. 2002. 秦岭泥盆系硅岩岩石类型对比研究. 吉林大学学报, 32(4): 325~328.

杨瑞东, 朱立军, 高慧, 张位华, 姜立君, 王强, 鲍森. 2005. 贵州遵义松林寒武系底部热液喷口及与喷口相关生物群特征. 地质论评, 51(5): 481~492.

杨瑞东, 魏怀瑞, 鲍森, 王伟, 王强. 2007. 贵州天柱上公塘—大河边寒武纪重晶石矿床海底热水喷流沉积结构, 构造特征. 地质论评, 53(5): 675~680.

杨水源, 姚静. 2008. 安徽巢湖平顶山中二叠统孤峰组硅质岩的地球化学特征及成因. 高校地质学报, 14(1): 39~48.

杨兴莲, 朱茂炎, 赵元龙, 张俊明, 郭庆军, 李丙霞. 2007. 黔东南前寒武纪—寒武纪转换时期微量元素地球化学特征研究. 地质学报, 81(10): 1391~1397.

杨兴莲, 赵元龙, 朱茂炎, 崔滔, 杨凯迪. 2010. 贵州丹寨寒武系牛蹄塘组海绵动物化石及其环境背景. 古生物学报, 49(3): 348~359.

杨玉卿, 冯增昭. 1997. 华南下二叠统层状硅岩的形成及意义. 岩石学报, 13(1): 111~120.

- 伊海生, 曾允孚, 夏文杰. 1989. 湘黔贵地区上震旦统沉积相及硅质岩成因研究. *矿物岩石*, 9(4): 54~58.
- 张春明, 张维生, 郭英海. 2012. 川东南—黔北地区龙马溪组沉积环境及对烃源岩的影响. *地学前缘*, 19(1): 136~145.
- 张位华, 姜立君, 高慧, 杨瑞东. 2003. 贵州寒武系底部黑色硅质岩成因及沉积环境探讨. *矿物岩石地球化学通报*, 22(2): 174~178.
- 张亚冠, 杜远生, 徐亚军, 余文超, 黄虎, 焦良轩. 2015. 湘中震旦纪—寒武纪之交硅质岩地球化学特征及成因环境研究. *地质论评*, 61(3): 499~510.
- 赵佩, 李贤庆, 田兴旺, 苏桂萍, 张明扬, 郭曼, 董泽亮, 孙萌萌, 王飞宇. 2014. 川南地区龙马溪组页岩气储层微孔隙结构特征. *天然气地球科学*, 25(6): 947~956.
- 周永章. 1990. 丹池盆地热水成因硅质岩地球化学特征. *沉积学报*, 8(3): 75~83.
- 周永章, 涂光炽, Edward H C, 卢焕章. 1994. 粤西古水剖面震旦系顶部层状硅岩的热水成因属性: 岩石学和地球化学证据. *沉积学报*, 12(3): 1~10.
- 周泽, 亢韦, 熊孟辉, 任海鹰, 闫高原. 2016. 贵州凤冈地区牛蹄塘组页岩气储层特征及勘探前景. *中国煤炭地质*, 28(6): 28~34.
- 朱嗣昭, Hiromi N, Masao K, 吴春寅. 1999. 一种特殊类型硅质岩—浮石状燧石. *高校地质学报*, 5(3): 345~350.
- Adachi M, Yamamoto K, Suigiski R. 1986. Hydrothermal chert and associated silicons rocks from the Northern Pacific: Their geological significance as indication of ocean ridge activity. *Sedimentary Geology*, 47: 125~148.
- Bi He, Jiang Zhenxue, Li Peng, Li Zhuo, Tang Xianglu, Zhang Dingyu, Xu Ye. 2014&. Shale reservoir characteristics and its influence on gas content of Wufeng—Longmaxi Formation in the southeastern Chongqing. *Natural Gas Geoscience*, 25(8): 1275~1283.
- Boggs J S. 2006. Application of Cathodoluminescence Imaging to the Study of Sedimentary Rocks. Cambridge University Press (177).
- Bostrom K, Peterson M N A. 1969. The origin of Al-poor ferromagnesian sediments in areas of high heat flow on the East Pacific Rise. *Geology*, 7: 427~447.
- Bostrom K. 1983. Genesis of Ferromanganese Deposits—Diagnostic Criteria for Recent and Old Deposits. New York: Plenum Press: 473~483.
- Calvert S E. 1983. Sedimentary geochemistry of silicon. In: Aston S R, ed. *Silicon Geochemistry and Biogeochemistry*: 143~170.
- Chen Daizhao, Wang Jianguo, Qing hairuo, Yan Detian, Li Renwei. 2009. Hydrothermal venting activities in the Early Cambrian, South China: Petrological, geochronological and stable isotopic constraints. *Chemical Geology*, 258: 168~181.
- Chen Hongde, Zeng Yunfu. 1989#. Discussion on the sedimentary characteristics and genesis of siliceous rocks of the Upper Devonian Liujiang Formation in Danchi Basin, Guangxi. *Mineral Rocks*, 9(4): 22~28.
- Clayton R N. 1986. High Temperature Isotope Effects in the Early Solar System. Houston: Pergamon Press: 1~139.
- Day-Stirrat R J, Milliken K L, Dutton S P, Loucks R G, Hillier S, Aplin A C, Schleicher A M. 2010. Open-system chemical behavior in deep Wilcox Group mudstones, Texas Gulf Coast, USA. *Marine and Petroleum Geology*, 27(9): 1804~1818.
- Dick J G. 2019. The microbiomes of deep-sea hydrothermal vents: Distributed globally, shaped locally. *Nature Reviews Microbiology*, 17(5): 271~283.
- Douthitt C B. 1982. The geochemistry of the stable isotopes of silicon. *Geochim Cosmochim Acta*, 46(8): 1449~1458.
- Fan Junxuan, Wu Lei, Chen Zhongyang, Chen Qing, Zhang Linna, Soo K Y, Singh N, Li Xuekang, Wang Yuan. 2013&. Biostratigraphy of the Wufeng—Longmaxi black shales at QiLin, Xingwen County, Sichuan Province. *Journal of Stratigraphy*, 37(4): 513~520.
- Fan Junxuan, Melchin M J, Chen Xu, Wang Yi, Zhang Yuandong, Chen Qing, Chi Zhaoli, Chen Feng. 2012#. Biostratigraphy of the black graptolite shale in the Ordovician—Silurian Longmaxi Formation in South China. *Science in China: Earth Sciences*, 42(1): 130~139.
- Feng Dongjun, Hu Zongquan, Gao Bo, Peng Yongmin, Du Wei. 2016&. Analysis of shale gas reservoir—forming condition of Wufeng Formation —Longmaxi Formation in southeast Sichuan Basin. *Geological Review*, 62(6): 1521~1532.
- Feng Shengbin, Zhou Hongrui, Yan Changhai, Peng Yi, Yuan Xiaoqi, He Jing. 2007&. Geochemical characteristics of hydrothermal cherts of Erlangping Group in east Qinling and their geologic significance. *Acta Sedimentologica Sinica*, 25(2): 564~573.
- Feng Mingyou, Zheng Jiang, Liu Tian, Liu Xiaohong, Hang Jiaqiang, Mei Wenhua, Sun Niyuan, Long Tao. 2018&. Characteristics and implications of fracture filling in carbonate concretions from the Lower Cambrian Qiongzhusi Formation, Northern Sichuan. *Geological Review*, 64(3): 711~722.
- Guidry S A, Chafetz H S. 2003. Anatomy of siliceous hot springs: examples from Yellowstone National Park, Wyoming, USA. *Sedimentary Geology*, 157(1~2): 71~106.
- Guo Fusheng, Lin Ziyu, Du Yangsong, Liu Linqing, Yan Zhaobin. 2003#. Research on the characteristics and genesis of a special type of siliceous rock. *Geoscience Frontiers*, 10(4): 573~581.
- Guo Tonglou, Zhang Hanrong. 2014&. Formation and enrichment mode of Jiaoshiba shale gas field, Sichuan Basin. *Petroleum Exploration and Development*, 41(1): 28~36.
- Han Shanchu, Hu Kai, Cao Jian. 2013&. First discovery of zoned hyalophane in the barite deposits hosted in Early Cambrian black shales of South China and its geological implications. *Geological Review*, 59(6): 1143~1149.
- Holdaway H K, Clayton C J. 1982. Preservation of shell microstructure in silicified brachiopods from the Upper Cretaceous Wilmington sands of Devon. *Geological Magazine*, 119: 371~382.
- Holland H D. 1967. Gangue minerals in hydrothermal system. In: Barners H. L., ed. *Geochemistry of Hydrothermal Ore Deposits*: 382~436.
- Hutton A C. 1994. Organic petrography and oil shales. Center for Applied Energy Research, 5(5): 1~4.
- Jarvie D M, Hill R J, Ruble T E, Pollastro R M. 2007. Unconventional shale—gas systems: The Mississippian Barnett Shale of north—central Texas as one model for thermogenic shale—gas assessment. *AAPG Bull*, 91: 475~499.
- Jiang Shaoyong, Yang Jinghong, Ling Hongfei, Feng Hongzhen, Chen Yongquan, Chen Jianhua. 2003. Re-Os isotopes and PGE geochemistry of black shales and intercalated Ni—Mo polymetallic sulfide bed from the Lower Cambrian Niutitang Formation, South China. *Progress in Natural Science*, 14(10): 788~794.
- Jin Zhijun, Hu Zongquan, Gao Bo, Zhao Jianhua. 2016&. Controlling factors on the enrichment and high productivity of shale gas in the Wufeng—Longmaxi Formations, southeastern Sichuan Basin. *Geoscience Frontiers*, 23(1): 1~9.
- Kamber B S, Webb G E. 2001. The geochemistry of late archaean microbial carbonate: Implications for ocean chemistry and continental erosion history. *Geochimica et Cosmochimica Acta*, 65:

2509~2525.

- Kenneth M H. 2005. Molecular phylogeny of siboglinid annelids (a. k. a. pogonorans): a review. *Hydrobiologia*, 535~536(1): 297~307.
- Krauskopf K B. 1956. Factors controlling the concentration of thirteen rare metals in sea-waters. *Geochemistry and Cosmochemistry Acta*, 9(1): 32.
- Leng Jigao, Gong Dajian, Li Fei, Li Peng. 2016. Analyses on the shale gas exploration prospect of the Niutitang Formation in the northeastern Guizhou area. *Earth Science Frontiers*, 23: 29~38.
- Li Ang, Ding Wenlong, Wang Ruyue, He Jianhua, Wang Xinghua, Sun Yaxiong, Gu Yang, Jiao Nailin. 2017. Petrophysical characterization of shale reservoir based on nuclear magnetic resonance (NMR) experiment: a case study of Lower Cambrian Qiongzhusi Formation in eastern Yunnan Province, South China. *Journal of Natural Gas Science and Engineering*, 37: 29~38.
- Li Fengjie, Liu Dianhe, Liu Qi. 2010. Geochemical characteristics and genesis discussion of Wujiaping Formation siliceous rocks in Xuanhan area of Sichuan Basin. *Natural Gas Geoscience*, 21(1): 62~67.
- Li Shengrong, Gao Zhenmin. 1995. Rare earth characteristics of the black rock series of the Niutitang Formation in the Xiangqian area; Also on the rare earth model of marine hydrothermal sedimentary rocks. *Marine Petroleum Geology*, 14(2): 1~19.
- Li Wenheng. 1980. A preliminary discussion on the genesis of a siliceous rock. *Geological Review*, 26(5): 442~445.
- Liang Digang, Guo Tonglou, Bian Lizeng, Chen Jianping, Zhao Feng. 2009. Controlling factors on the sedimentary facies and development of Palaeozoic marine source rocks. *Marine Petroleum Geology*, 14(2): 1~19.
- Liu Honglin, Wang Hongyan, Fang Chaohe, Guo Wei, Sun Shasha. 2016. The formation mechanism of over-pressure reservoir and target screening index of the marine shale in the South China. *Earth Science Frontiers*, 23: 48~54.
- Liu Jiangtao, Li Yongjie, Zhang Yuanchun, Liu Shuanglian, Cai Yingjie. 2017. Evidences of biogenic silica of Wufeng—Longmaxi Formation shale in Jiaoshiba area and its geological significance. *Journal of China University of Petroleum (Edition of Natural Science)*, 41(1): 34~41.
- Liu Shugen, Ma Wenxin, Luba J, Huang Wen, Zeng Xiangliang, Zhang Changjun. 2011. Characteristics of the shale gas reservoir rocks in the Lower Silurian Longmaxi Formation, east Sichuan basin, China. *Acta Petrologica Sinica*, 27(8): 2239~2252.
- Liu Shugen, Wang Shiyu, Sun Wei, Ran Bo, Yang Di, Luo Chao, Ye Yuehao, Bai Zhiqiang, Qiu Jiawen. 2013. Characteristics of black shale in Wufeng Formation and Longmaxi Formation in Sichuan Basin and its peripheral areas. *Journal of Chengdu University of Technology: Natural Science Edition*, 40(6): 621~639.
- Liu Wei, Xu Xiaosong, Feng Xintao, Sun Yuanyuan. 2010. Radiolarian siliceous rocks and palaeoenvironmental reconstruction for the Upper Ordovician Wufeng Formation in the middle—upper Yangtze area. *Sedimentation and Tethyan Geology*, 30(3): 65~70.
- Liu Z H, Zhuang X G, Teng G E, Xie X M, Yin L M, Bian L Z, Feng Q L, Algeo T J. 2015. The lower Cambrian Niutitang Formation at Yangtiao (Guizhou, SW China): Organic matter enrichment, source rock potential, and hydrothermal influences. *Journal of Petroleum Geology*, 38(4): 11~432.
- Loucks R G, Ruppel S C. 2007. Mississippian Barnett Shale: Lithofacies and depositional setting of a deep-water shale-gas succession in the Fort Worth basin, Texas. *AAPG Bull*, 91: 579~601.
- Lu Longfei, Qin Jianzhong, Shen Baojian, Tenggeer, Liu Weixin, Zhang Qingzhen. 2016. Biogenic origin and hydrocarbon significance of siliceous shale from the Wufeng—Longmaxi formations in Fuling area, southeastern Sichuan Basin. *Petroleum Geology and Experiment*, 38(4): 460~466.
- Marin-Carbonne J, Chaussidon M, Boiron M C, Robert F. 2011. A combined in situ oxygen, silicon isotopic and fluid inclusion study of a chert sample from Onverwacht Group (3.35 Ga, South Africa): New constraints on fluid circulation. *Chemical Geology*, 286: 59~71.
- Milliken K L, Esch W L, Reed R M, Zhang T W. 2012. Grain assemblages and strong diagenetic overprinting in siliceous mudrocks, Barnett Shale (Mississippian), Fort Worth Basin, Texas. *AAPG Bull*, 96: 1553~1578.
- Murray R W. 1990. Rare earth element as indicators of different marine depositional environments in chert and shale. *Geology*, 18: 268~271.
- Murray R W, Tenbrink M R B, Gerlach D C. 1991. Rare earth, major, and trace elements in chert from the Franciscan complex and Monterey group, Californian: Assessing REE sources to fine-grained marine sediments. *Geochimica et Cosmochimica Acta*, 55: 1875~1895.
- Murray R W. 1992. Interoceanic variation in the rare earth, major and trace element depositional chemistry of chert: Perspectives gained from DSDP and ODP record. *Geochim Cosmochim Acta*, 56: 1897~1913.
- Monika B, Olav G. 2005. Microbial Symbiosis in Annelida. *Symbiosis*, 38: 1~45.
- Rakociński M, Boreczek E. 2016. Concentrations of silicified cephalopods within upper Frasnian carbonate concretions from the Holy Cross Mountains, Poland. *Palaeogeography Palaeoclimatology Palaeoecology*, 449: 475~483.
- Robert G L, Stephen C R. 2007. Mississippian Barnett Shale: Lithofacies and depositional setting of a deep-water shale-gas succession in the Fort Worth Basin, Texas. *AAPG Bulletin*, 91(4): 579~601.
- Rona P A. 1988. Hydrothermal mineralization of ocean ridges. *Canadian Mineralogy*, 26(3): 447~465.
- Rona P A. 1978. Criteria for recognition of hydrothermal mineral deposits in ocean crust. *Economic Geology*, 73: 135~160.
- Rona P A, Bostrom K, Epstein S. 1980. Hydrothermal quartz vug from the Mid-Atlantic Ridge. *Geology*, 8: 569~572.
- Schieber J, Krinsley D, Riciputi L. 2000. Diagenetic origin of quartz silt in mudstones and implications for silica cycling. *Nature*, 406: 981~985.
- Shao Xinhe, Pang Xiongqi, Li Hui, Zhang Xue. 2017. Fractal analysis of pore network in tight gas sandstones using NMR method: A case study from the Ordos Basin, China. *Energy Fuel*, 31(10): 10358~10368.
- Sheng Jihu, Chen Zhongxin. 1998. Research status of siliceous rock sedimentary geochemistry. *Regional Research and Development*, 17(Supplement): 117~121.
- Shi Guijun, Yang Xiangning. 1999. Siliceous limestone of Huanglong Formation in the junction area of southern Anhui and western Zhejiang. *Geological Journal of China Universities*, 5(1): 113~114.
- Song Tianrui, Ding Tiping. 1989. A new attempt to apply the silicon

- isotope ($\delta^{30}\text{Si}$) in siliceous rock to sedimentary facies analysis. *Chinese Science Bulletin*, 34(18):1408~1411.
- Steiner M, Wallis E, Erdtmann B D, Zhao Y L, Yang R D. 2001. Submarine-hydrothermal exhalative ore layers in black shales from South China and associated fossils—insights into a Lower Cambrian facies and bio-evolution. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 169: 165~191.
- Sun Wei, Liu Shugen, Ran Bo, Wang Shiyu, Ye Yuehao, Luo Chao, Tian Mengna. 2012&. General situation and prospect evaluation of the shale gas in Niutitang Formation of Sichuan Basin and its surrounding areas. *Journal of Chengdu University of Technology (Natural Science Edition)*, 39(2): 170~175.
- Tan Jingqiang, Horsfield B, Fink R, Krooss B, Schulz H M, Rybacki E, Zhang Jinchuan, Boreham C J, Graas G V, Tocher B A. 2014. Shale gas potential of the major marine shale formations in the upper Yangtze platform, South China, Part III: Mineralogical, lithofacial, petrophysical, and rock mechanical properties. *Energy Fuels* 28(4): 2322~2342.
- Tian Yuntao, Feng Qinglai, Li Qin. 2007&. The petrogenesis and sedimentary environment of the bedded cherts from Upper Permian Dalong Formation, southwest Guangxi. *Acta Sedimentologica Sinica*, 25(5): 671~677.
- Tu Guangchi. 1990#. Progress in several issues in earth sciences in the 1980s. *Geological Review*, 36(6): 510~517.
- Wang Chao, Zhang Baiqiao, Lu Yongchao, Shu Zhiguo, Lu Yaqiu, Bao Hanyong, Meng Zhiyong, Chen Lei. 2018. Lithofacies distribution characteristics and its controlling factors of shale in Wufeng Formation—Member 1 of Longmaxi Formation in the Jiaoshiba area. *Petroleum Research* 3: 306~319.
- Wang Ruyue, Gong Dajian, Ding Wenlong, Leng Jigao, Yin Shuai, Wang Xinghua, Sun Yaxiong. 2016. Brittleness evaluation of the Lower Cambrian Niutitang formation shale in the upper Yangtze region; A case study in the Cengong block, Guizhou province. *Earth Science Frontiers*, 23: 87~95.
- Wang Shufang, Zou Caineng, Dong Dazhong, Wang Yuman, Huang Jinliang, Guo Zhaojie. 2014&. Biogenic silica of organic-rich shale in Sichuan Basin and its significance for shale gas. *Journal of Peking University: Natural Science Edition*, 50(3): 476~486.
- Wang Yuman, Dong Dazhong, Li Jianzhong, Wang Shejiao, Li Xinjing, Wang Li, Cheng Keming, Huang Jinliang. 2012&. Reservoir characteristics of shale gas in Longmaxi Formation of the Lower Silurian, southern Sichuan. *Acta Petrolei Sinica*, 33(4): 551~561.
- Wang Yuman, Dong Dazhong, Li Xinjing, Huang Jinliang, Wang Shufang, Wu Wei. 2015&. Stratigraphic sequence and sedimentary characteristics of Lower Silurian Longmaxi Formation in the Sichuan Basin and its peripheral areas. *Natural Gas Industry*, 35(3): 12~21.
- Wang Yuman, Wang Shufang, Dong Dazhong, Li Xinchun, Huang Jinliang, Zhang Chenchen, Guan Quanzhong. 2016&. Lithofacies characterization of Longmaxi Formation of the Lower Silurian, southern Sichuan. *Geoscience Frontiers*, 23(1): 119~133.
- Wei Xiangfeng, Zhao Zhengbao, Wang Qingbo, Liu Zhujiang, Zhou Min, Zhanghui. 2017&. Comprehensive evaluation on geological conditions of the shale gas in Upper Ordovician Wufeng Formation—Lower Silurian Longmaxi Formation in Dingshan area, Qijiang, Southeastern Sichuan. *Geological Review*, 63(1): 153~164.
- Wu Anbin, Zhang Jingkun, Wang Jingling, Luo Jiaguo, Luo Qun, Jiang Zhenxue. 2020&. Genesis, diagenetic model and geological significance of calcite veins in organic rich shale; a case study of the Longmaxi Formation, southern Sichuan basin, China. *Geological Review*, 66(1): 88~99.
- Wu Lanyu, Hu Dongfeng, Lu Yongchao, Liu Ruobing, Liu Xiaofeng. 2016&. Advantageous shale lithofacies of Wufeng Formation—Longmaxi Formation in fuling gas field of Sichuan Basin, SW China. *Petroleum Exploration and Development*, 43(2): 189~197.
- Xia Bangdong, Zhong Lirong, Fang Zhong, Lv Hongbo. 1995#. The genesis of the layered siliceous rocks in the Gufeng Formation of the Early Permian in the lower Yangtze district. *Acta Geology*, 69(2): 125~137.
- Xie Xiaomin, Teng Geer, Qin Jianzhong, Zhang Qingzhen, Bian Lizeng, Yin Leiming. 2015&. Depositional environment, organisms components and source rock formation of siliceous rocks in the base of the Cambrian Niutitang Formation, Kaili, Guizhou. *Acta Geology*, 89(2): 425~439.
- Xu Zhuang, Shi Wanzhong, Zhai Gangyi, Clay C, Zhang Xiaoming, Peng Nyujia. 2019. A rock physics model for characterizing the total porosity and velocity of shale: A case study in Fuling area, China. *Marine and Petroleum Geology*, 99: 208~226.
- Yamamoto K. 1987. Geochemical characteristics and depositional environment of cherts and associated rocks in the Franciscan and Shimanto terraces. *Sedimentary Geology*, 52: 65~108.
- Yang Feng, Ning Zhengfu, Wang Qing, Zhang Rui, Krooss B M. 2016. Pore structure characteristics of lower Silurian shales in the southern Sichuan Basin, China: Insights to pore development and gas storage mechanism. *International Journal of Coal Geology* 156: 12~24.
- Yang Jian, Yi Fazheng, Hou Lanjie. 2004&. Genesis and petrogeochemistry characteristics of Lower Cambrian black shales series in Northern Guizhou. *Acta Minera Sinica*, 24(3): 285~289.
- Yang Ji, Li Ying. 2002&. Discussion of silica rock types by comparsion in devonian system of Qinling Mountains. *Journal of Jilin University*, 32(4): 325~328.
- Yang Ruidong, Zhu Lijun, Gao Hui, Zhang Weihua, Jiang Lijun, Wang Qiang, Bao Miao. 2005#. The characteristics of hydrothermal vents at the bottom of the Cambrian in the pine forests of Zunyi, Guizhou and related biota. *Geological Review*, 51(5): 481~492.
- Yang Ruidong, Wei Huairui, Bao Miao, Wang Wei, Wang Qiang. 2007&. Submarine hydrothermal venting—flowing sedimentary characters of the Cambrian Shanggongtang and Dahebian barite deposits, Tianzhu County, Guizhou Province. *Geological Review*, 53(5): 675~680.
- Yang Shuiyuan, Yao Jing. 2008&. Geochemistry and origin of siliceous rocks from the Gufeng Formation of Middle Permian in the Pingdingshan Area, Chaohu Region, Anhui Province. *Geological Journal of China Universities*, 14(1): 39~48.
- Yang Xiangrong, Yan Detian, Wei Xiaosong, Zhang Liwei, Zhang Bao, Xu Hanwen, Gong Yin, He Jie. 2018. Different formation mechanism of quartz in siliceous and argillaceous shales; A case study of Longmaxi Formation in South China. *Marine and Petroleum Geology*, 94: 80~94.
- Yang Xinglian, Zhu Maoyan, Zhao Yuanlong, Zhang Junming, Guo Qingjun, Li Bingxia. 2007&. Trace element geochemical characteristics from the Ediacaran Cambrian Transition interval in eastern Guizhou, South China. *Chinese Journal of Geology*, 81(10): 1391~1397.
- Yang Xinglian, Zhao Yuanlong, Zhu Maoyan, Cui Tao, Yang Kaidi. 2010&. Sponges from the Early Cambrian Niutitang Formation at Danzhai, Guizhou and their environmental background. *Chinese Journal of Paleontology*, 49(3): 348~359.

Yang Yuqing, Feng Zengzhao. 1997#. The formation and significance of the Lower Permian layered silica in South China. *Acta Petrologica Sinica*, 13(1): 111~120.

Yi Haisheng, Zeng Yunfu, Xia Wenjie. 1989#. Study on the sedimentary facies and siliceous origin of the Upper Sinian in the Xiangqiangui area. *Mineralogy*, 9(4): 54~58.

Zeng Weite, Ding Wenlong, Zhang Jinchuan, Li Yuxi, Lin Tuo, Wang Ruyue, Fu Jinglong. 2016. Research on the fracture effectiveness of the lower cambrian Niutitang shale in the southeastern Chongqing and northern Guizhou areas. *Earth Sci. Frontiers*, 23: 96~106.

Zhang Chunming, Zhang Weisheng, Guo Yinghai. 2012#. Sedimentary environment and its effect on hydrocarbon source rocks of Longmaxi Formation in southeast Sichuan and northern Guizhou. *Geoscience Frontiers*, 19(1): 136~145.

Zhang Kun, Song Yan, Jiang Shu, Jiang Zhenxue, Jia Chengzao, Huang Yizhou, Wen Ming, Liu Weiwei, Xie Xuelian, Liu Tianlin, Wang Pengfei, Shan Changan, Wu Yinghui. 2018. Mechanism analysis of organic matter enrichment in different sedimentary backgrounds: A case study of the Lower Cambrian and the Upper Ordovician—Lower Silurian, in Yangtze region. *Marine and Petroleum Geology*, 99: 488~497.

Zhang Weihua, Jiang Lijun, Gao Hui, Yang Ruidong. 2003#. Study on sedimentary environment and origin of black siliceous rocks of the Lower Cambrian in Guizhou Province. *Petrology and Geochemistry*, 22(2): 174~178.

Zhang Yaguan, Du Yuansheng, Xu Yajun, Yu Wenchao, Huang Hu, Jiao Liangxuan. 2015#. Geochemical characteristics of siliceous rocks during the transition from Sinian (Ediacaran) to Cambrian in Central Hunan and its implication for genesis and sedimentary environment. *Geological Review*, 61(3): 499~510.

Zhang Yuying, He Zhiliang, Jiang Shu, Lu Shuangfang, Xiao Dianshi, Chen Guohui, Zhao Jianhua. 2018. Factors affecting shale gas accumulation in the overmature shales——case study from the lower Cambrian shale in the western Sichuan Basin, South China. *Energy Fuels*, 32: 3003~3012.

Zhang Yuying, He Zhiliang, Jiang Shu, Lu Shuangfang, Xiao Dianshi, Chen Guohui, Li Yuchao. 2019. Fracture types in the lower Cambrian shale and their effect on shale gas accumulation, Upper Yangtze. *Marine and Petroleum Geology*, 99: 282~291.

Zhao Jianhua, Jin Zhijun, Hu Qinhong, Liu Keyu, Jin Zhenkui, Hu Zongquan, Nie Haikuan, Du Wei, Yan Caina, Wang Ruyue. 2018. Mineral composition and seal condition implicated in pore structure development of organic-rich Longmaxi shales, Sichuan Basin, China. *Marine and Petroleum Geology*, 98: 507~522.

Zhao Pei, Li Xianqing, Tian Xingwang, Su Guiping, Zhang Mingyang, Guo Man, Dong Zeliang, Sun Mengmeng, Wang Feiyu. 2014#. Study on micropore structure characteristics of Longmaxi Formation shale gas reservoirs in the southern Sichuan Basin. *Natural Gas Geoscience*, 25(6): 947~956.

Zhou Yongzhang. 1990#. Geochemical characteristics of siliceous rocks of hydrothermal origin in Danchi Basin. *Acta Sedimentologica Sinica*, 8(3): 75~83.

Zhou Yongzhang, Tu Guangchi, Edward H C, Lu Huanzhang. 1994#. Hydrothermal genetic properties of the top Sinian layered silica at the Gushui section in western Guangdong: evidence from petrology and geochemistry. *Acta Sedimentologica Sinica*, 12(3): 1~10.

Zhou Ze, Kang Wei, Xiong Menghui, Ren Haiying, Yan Gaoyuan. 2016#. Niutitang Formation shale gas reservoir features and exploration prospect in Fenggang Area, Guizhou. *China Coal Geology*, 28(6): 28~34.

Zhu Sizhao, Hiromi N, Masao K, Wu Chunyin. 1999#. A special type of siliceous rock—pumice-like flint. *Geological Journal of China Universities*, 5(3): 345~350.

Siliceous source and its influence on organic matter preservation of the two Lower Paleozoic siliceous shale reserviors in Sichuan Basin

XIE Xiaomin^{1, 2, 3)}, LIU Weixin³⁾, ZHANG Yin⁴⁾, ZHAO Difei⁵⁾,
TANG Youjun¹⁾, SHEN Baojian³⁾, LI Zhiming³⁾

- 1) *Key Laboratory of Oil and Gas Resources and Exploration Technology, Ministry of Education, College of Resources and Environment, Yangtze University, Wuhan, 430100;*
- 2) *Nanjing Hongchuang Exploration Technology Service Co. Ltd, Nanjing, 211111;*
- 3) *Wuxi Institute of Petroleum Geology, SINOPEC, Wuxi, Jiangsu, 214126;*
- 4) *Chongqing University of Science and Technology, Chongqing, 401331;*
- 5) *China University of Mining and Technology, Xuzhou, Jiangsu, 221008*

Abstract: Two sets of siliceous shale formations in Lower Paleozoic, the bottom of Cambrian and the Ordovician—Silurian Wufeng—Longmaxi Shales, are the important shale reservoirs in Sichuan Basin. The TOC and shale gas content of the Wufeng—Longmaxi shales are positively related to the siliceous quantity, suggesting that the siliceous materials could be very important for the shale gas formation. However, the origin of siliceous materials is complicated, and there could be some siliceous fluid filled after the rock sedimentation. This study, based on the summary of previous research results, has carefully analyzed the petrographical characteristics of siliceous minerals. Three types of siliceous sources have been detected: ① syngenetic hydrothermal inorganic siliceous fluid, ② bio-silica, and ③ late-filled inorganic siliceous fluid. Syngenetic hydrothermal inorganic siliceous fluid and bio-silica are the dominated siliceous sources in Lower Cambrian shales. Syngenetic hydrothermal inorganic siliceous fluid made the organic matter rapidly silicified, and the organic matter could be

exquisitely preserved. The siliceous source in the Wufeng—Longmaxi Formation was mainly from late-filled inorganic siliceous fluid and bio-silica, and organic matter preservation has little influence by siliceous fluid. Thus, the organic matter has gone through slowly saprofitication process, resulting in the morphology of organic matters are poor preserved. The study has initially revealed the petrographical characteristics of siliceous materials and the influences on the organism preservations for comparision the Lower Cambrian and the Wufeng—Longmaxi shale reservoirs, with hoping to provide new geological data and ideas for shale gas exploration in Sichuan Basin.

Keywords: siliceous shale; origin of siliceous materials; organic matter preservation; Lower Paleozoic; Sichuan Basin

Acknowledgement: This research was supported by the National Natural Science Foundation of China (No. 41972163)

First author: XIE Xiaomin, female, born in 1984, Ph. D. , is now focus on the combination researches of petrology and geochemistry in Yangtze University; Email: xiaominxie2019@sina.com

Manuscript received on: 2020-06-14; Accepted on: 2021-01-11; Network published on: 2021-02-20

Doi: 10. 16509/j. georeview. 2021. 02. 012 **Edited by:** LIU Zhiqiang

中国地质学会 2020 年度“十大地质找矿成果”

(按矿藏所在省、市、自治区名汉语拼音字母为序)

1 重庆东南南川招标区探明超千亿方大型常压页岩气田

中国石化华东油气分公司历经十余年攻关,在渝东南南川招标区发现了我国首个超千亿方大型常压页岩气田。针对常压页岩气富集规律复杂、甜点优选难度大、单井产量和EUR(估算最终采收量)低等难点,攻关团队创新提出了“沉积相带、保存条件、地应力场”三因素控产地质理论,构建了高陡背斜、反向逆断层遮挡等四种成藏模式,创建了“岩相-应力-孔隙”一体化储层综合评价、变密度三维地震勘探、基于地质-工程一体化的甜点评价及水平井设计技术等勘探关键技术,实现勘探重大突破,提交探明储量 $1918\times10^8\text{ m}^3$ 。南川常压页岩气田的发现是我国页岩气地质理论创新和勘探技术突破的重大成果,有效推动了我国南方 $9\times10^{12}\text{ m}^3$ 常压页岩气资源的效益开发。

2 江西省上饶市许家桥发现世界最大滑石矿

以江西省地质勘查基金管理中心王先广为首席的科研团队,在江西省上饶市广丰区许家桥矿区探获世界资源储量规模最大的滑石矿,提交(控制+推断)滑石资源量 95. 523 Mt(相当全国新增滑石 32%),其中控制滑石矿资源量 40. 367 Mt。上饶市广丰区许家桥滑石矿的发现,刷新了世界滑石矿资源储量规模纪录,改变了世界滑石矿分布格局;创建了“同生断层+富硅热液+富镁海水+有机质”黑滑石成矿模型和“许家桥式”滑石矿“层位+钙镁硅质岩石组合+同生断层+向斜构造”找矿模式,深化了区域成矿规律。区内黑滑石矿石品质好,创建的黑滑石晶体结构模型开拓了材料应用新方向,为上饶市打造千亿级滑石矿产业集群提供了资源保障。

3 内蒙古阿拉善右旗塔木素矿区发现超大型天然碱矿

内蒙古矿业开发有限责任公司在阿拉善右旗塔木素矿区取得天然碱矿重大找矿突破,获得固体天然碱(122b+333)矿石量 1. 078 Gt,矿物量($\text{Na}_2\text{CO}_3+\text{NaHCO}_3$) 709. 09 Mt,其中控制的经济基础储量(122b) 天然碱矿石量 608. 84 Mt,矿物量 406. 33 Mt,是我国目前已查明天然碱储量最大的天然碱矿。其成矿时代为白垩纪,突破了我国天然碱矿仅在新近纪以来成矿的历史,为今后天然碱矿的发现提供了更大的找矿空间。该天然碱矿的发现,或将改变我国以化学合成法制碱为主的现状,减轻化学制碱对自然环境造成的压力,对环境保护起到积极作用,并将带来较大的社会经济效益。

4 青海省都兰县那更康切尔沟探明超大型银多金属矿

由四川省冶金地质勘查局水文工程大队承担的“青海省都兰县那更康切尔沟银多金属普查”项目取得重大找矿突破,在青海东昆仑地区首次发现了超大型中低温构造热液型银矿床,地质勘查、研究意义重大。团队通过地质、物探、化探、科研等多专业协同合作,产学研联合攻关,在矿区内已发现银矿带 18 条,圈出银矿体 69 条,评审认定(333+334?)银资源量 5070. 56 t,矿床平均品位 $326. 35\times10^{-6}$,其中(333)资源量 2137. 2 t,伴生铅锌 0. 152 Mt,深部及外围还具有较大的找矿潜力。那更康切尔沟银矿的发现,填补了青海省无超大型独立银矿床的空白,对青海省的经济建设以及西部大开发都具有重要意义。

5 山西省繁峙县发现大型斑岩型金矿

紫金矿业集团股份有限公司地质找矿项目团队通过总结繁峙县义兴寨金矿床的成矿规律, (下转第 475 页)